

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number
WO 01/36674 A2

(51) International Patent Classification: C12Q 1/68,
G01N 33/53, A61P 43/00

Dora [IE/GB]; 45 St. Anne's Avenue, Grappenhall, War-
rington WA4 2PL (GB).

(21) International Application Number: PCT/GB00/04267

(74) Agent: GAINNEY, Laurence, David, Scott; Astrazeneca,
Global Intellectual Property, P.O. Box 272, Mereside,
Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).

(22) International Filing Date:
8 November 2000 (08.11.2000)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9926805.4 13 November 1999 (13.11.1999) GB

(71) Applicant (*for all designated States except MG, US*): AS-
TRAZENECA AB [SE/SE]; S-151 85 Sodertalje (SE).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for MG only*): ASTRAZENECA UK LIM-
ITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN
(GB).

Published:

— Without international search report and to be republished
upon receipt of that report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BULL, John, Henry
[GB/GB]; 25 Pownall Street, Macclesfield, Cheshire SK10
1DF (GB). ELLISON, Gillian [GB/GB]; Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). PASKINS, Lynn,

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: DIAGNOSTIC METHODS

(57) Abstract: The present invention derives from the identification of 26 gene transcripts (Markers) that exhibit aberrant expres-
sion levels in prostate disorder tissues. The invention therefore relates to diagnostic techniques for the detection of human prostate
disorders, such as cancer, by detecting one or more of these Markers, intermediates, precursors or products (mRNA, cDNA, genomic
DNA, or protein). The invention is also directed to methods for identifying modulators of prostate disorders, which modulators, such
as chemical compounds, antisense molecules and antibodies interact with and modulate any one of the Markers identified.

WO 01/36674 A2

PTS-0308

- 2 -

In early disease, where the cancer is confined to the prostate gland and has not spread beyond the gland, the tumour can potentially be removed by surgery. Surgery is a common treatment of prostate cancer. Transurethral resection is a procedure in which the cancer is cut from the prostate via the urethra. A radical prostatectomy is where the whole prostate and
5 some of the tissue around the gland is removed. Radiation therapy is also used to treat early prostate cancer.

Once disease has become metastatic, its course is rapid and relatively predictable, leading to death within 2-5 years if untreated.

Prostate cancer commonly occurs in men above the age of 50 years. The incidence of
10 prostate cancer rises sharply between the ages of 60 and 80 years, with more than 80% of all cases of the diseases being diagnosed in men over 65 and less than 1% in men under 50 years of age. The incidence and prevalence of prostate cancer have increased considerably throughout the world over the past two decades. The average age of detection is 72 in the UK and 66 in the US. Active screening programmes may lead to a lowering of the average age at
15 first diagnosis.

Routine medical examinations for obstructive urinary symptoms may lead to an initial diagnosis of prostate cancer. When prostate cancer is asymptomatic, there are two tests that can be performed to detect the presence of cancer: digital rectal examination (DRE) and prostate specific antigen (PSA). PSA is an enzyme which is secreted almost exclusively by
20 epithelial cells in the prostate; measurement of the quantity of PSA in the blood provides a more reliable indicator of the presence of prostate cancer than the DRE test. However, elevations in serum levels of PSA can also be found in patients with benign prostatic hyperplasia (BPH) and prostatitis. In some cases, where a patient has elevated levels of PSA in his blood, a third diagnostic technique is used called 'transrectal ultrasonography' (TRUS),
25 which can further substantiate the presence of prostate cancer. In conjunction with taking a biopsy of the tissue, TRUS has been shown to increase the detection rate of prostate cancer two-fold, when compared to DRE alone. The combination of digital rectal examination, serum PSA level and transrectal ultrasound is currently the best available diagnostic tool (Gorgoulis VG. et al. Anticancer Res 19:2327-2348 (1999)).

30 It is common for the prostate gland to become enlarged with ageing, a condition called benign prostatic hyperplasia (BPH), or benign prostatic hypertrophy. Severe BPH can cause

- 4 -

nucleic acid targeted to the Marker mRNA; or more widely in the identification or development of chemical or hormonal therapeutic agents. The person skilled in the art is also capable of devising screening assays to identify compounds (chemical or biological) that modulate (activate or inhibit) one or other of the identified Marker, which compounds may
5 prove useful as therapeutic agents in treating a prostate disorder. Monitoring the Markers could also be useful in identifying inhibitors, antagonists or biochemical signalling effects in high throughput screening.

According to a first aspect of the invention there is provided a method for diagnosing or prognosing or monitoring a prostate disorder comprising testing a biological sample for
10 aberrant levels of one or more of the Markers selected from the group consisting of Marker 1 to Marker 26. Preferred Markers are numbers: 8, 13, 14, 15, 17, 18, 19, 20, 22, 24, 25 and 26.

The invention lies in the identification of these differentially expressed Markers in prostate disorders. Accordingly, the invention is directed to any diagnostic method capable of
15 assessing the differential expression levels, relative to expression in control tissues, of one or more of the identified Markers, either alone or as a panel. In particular, such methods include assessment of mRNA transcript levels and/or protein levels. The presence of aberrant expression levels of one of the Markers indicating the presence of a prostate disorder.

In a preferred embodiment the diagnostic method involves testing for more than one of
20 the Markers identified herein. As separate independent embodiments the diagnostic method may involve testing for 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or all of the Markers identified herein, optionally as part of a panel test of other gene Markers.

It will be apparent to the person skilled in the art that there are a large number of
25 analytical procedures which may be used to detect the amount of any of the Marker products present in a test sample.

The test sample comprising nucleic acid or protein is conveniently any prostatic material or biological sample, including TURP chip, biopsy, excised prostate or part thereof, a sample of bone marrow aspirate, bone marrow biopsy, lymph node aspirate, lymph node
30 biopsy, spleen tissue, fine needle aspirate, skin biopsy or organ tissue biopsy, particularly prostate gland tissue, invaded surgical margin, invaded lymph node, invaded lung, invaded

- 6 -

biomarkers for distinguishing cancer from BPH. Highly specific yet low abundance Markers could be invaluable if secreted in protein form, as they could permit non-invasive testing of body fluids.

According to a further aspect of the invention there is provided a method for
5 distinguishing prostate cancer from BPH comprising testing a biological sample for aberrant levels of one or more of the Markers selected from the group consisting of Marker 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 17, 18, 20, 23, 25 and 26. Such a method might involve measuring the expression levels of one or more of these Markers in a biological sample taken from an individual, comparing the Marker expression level detected with control values or
10 historical values and depending on the value detected determining whether or not the individual has prostate cancer or BPH.

Each of the Marker genes useful in the present invention are already in the public domain. However, the inventors are not aware of any prior art disclosing an association of any of the Marker genes with prostate cancer. Thus, although each of the Markers/genes
15 identified herein are themselves already known, their association with prostate disorders leading to the present invention is unknown.

Marker 1 is the human smooth muscle protein, 22kDa. The gene coding for this protein has been cloned and sequenced (Thweatt et al., Biochem. Biophys. Res. Commun. 187:1-7 (1992)). These authors identify 22kD smooth muscle protein as a fibroblast or smooth
20 muscle protein. There is no teaching in this paper of an association with prostate cancer. The sequence of 22kDa smooth muscle protein is disclosed in Thweatt et al. and is present in the EMBL database under accession number HS22SM. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HS22SM, unless stated otherwise or apparent from
25 the context.

Marker 2 is the Rho GDP dissociation inhibitor protein. The gene coding for this protein has been cloned and sequenced (Leffers et al., Exp. Cell Res. 209:165-174 (1993)). These authors identify Rho GDP dissociation inhibitor protein as a keratinocyte protein. There is no teaching in this paper of an association with prostate cancer. The sequence of Rho
30 GDP dissociation inhibitor protein is disclosed in Leffers et al., and is present in the EMBL database under accession number HSRHO1. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL

- 8 -

acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSD966, unless stated otherwise or apparent from the context.

Marker 7 is the human mRNA KIAA0120. This cDNA has been cloned and sequenced (Nagase et al., DNA Res. 2:37-43 (1995)). These authors identify KIAA0120 as a
5 brain transcript. There is no teaching in this paper of an association with prostate cancer. The sequence KIAA0120 is disclosed in Nagase et al., and is present in the EMBL database under accession number HSORFF. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSORFF, unless stated otherwise or apparent from the context.

10 Marker 8 is the human ribosomal protein S25. The gene coding for this protein has been cloned and sequenced (Li et al., Gene 107:329-333 (1991)). These authors identify ribosomal protein S25 as a leukaemia cell line protein. There is no teaching in this paper of an association with prostate cancer. The sequence of human ribosomal protein S25 is disclosed in Li et al., and is present in the EMBL database under accession number HSRPS25.
15 For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSRPS25, unless stated otherwise or apparent from the context.

Marker 9 is the human 80K-H protein (kinase C substrate). The gene coding for this protein has been cloned and sequenced (Sakai et al., Genomics 5:309-315 (1989)). These
20 authors identify 80K-H protein (kinase C substrate) as a fibroblast and epidermal carcinoma cell protein. There is no teaching in this paper of an association with prostate cancer. The sequence of 80K-H protein (kinase C substrate) is disclosed in Sakai et al., and is present in the EMBL database under accession number HSG19P1A. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in
25 the EMBL database under accession number HSG19P1A, unless stated otherwise or apparent from the context.

Marker 10 is the human alpha-2-macroglobulin protease inhibitor protein. The gene coding for this protein has been cloned and sequenced (Kan et al., Proc. Natl. Acad. Sci. U.S.A. 82:2282-2286 (1985)). These authors identify alpha-2-macroglobulin protease
30 inhibitor protein as a serum protein. There is no teaching in this paper of an association with prostate cancer. The sequence of alpha-2-macroglobulin protease inhibitor protein is disclosed in Kan et al., and is present in the EMBL database under accession number

- 10 -

referred to herein, refers to that in the EMBL database under accession number HSLB2A26, unless stated otherwise or apparent from the context.

Marker 15 is the human PRSM1 protein. The gene for PRSM1 has been cloned and sequenced (Scott et al., Gene 174:135-143 (1996)). These authors identify PRSM1 as a putative metallopeptidase. There is no teaching in this paper of any association with prostate cancer. The sequence of PRSM1 is disclosed in Scott et al. and is present in the EMBL database under accession number HSU58048. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSU58048, unless stated otherwise or apparent from the context.

Marker 16 is the human ribosomal large subunit L12 protein. The gene for ribosomal large subunit L12 protein has been cloned and sequenced (Chu et al., Nucleic Acids Res. 21:749-749 (1993)). These authors identify human ribosomal large subunit L12 as a ribosomal protein. There is no teaching in this paper of any association with prostate cancer. The sequence of human ribosomal large subunit L12 is disclosed in Chu et al. and is present in the EMBL database under accession number HSL12A. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSL12A, unless stated otherwise or apparent from the context.

Marker 17 is the human 7SK45 RNA. The gene for 7SK45 RNA has been cloned and sequenced (Murphy et al., Nucleic Acids Res. 14:9243-9260 (1986)). These authors identify human 7SK45 as a small cytoplasmic RNA. There is no teaching in this paper of any association with prostate cancer. The sequence of 7SK45 is disclosed in Murphy et al. and is present in the EMBL database under accession number HS7K45. For the purpose of this application, the gene sequences and sequence positions referred to herein, refers to that in the EMBL database under accession numbers: HS7K45, unless stated otherwise or apparent from the context.

Marker 18 is the human KIAA0588 protein (human protocadherin gamma). This protein is abundantly expressed in brain cells. The gene for this protein has been cloned and sequenced (Nagase et al., DNA Research. 5:31-39 (1998)). There is no teaching in this paper of an association with prostate cancer. The sequence of the KIAA0588 protein is present in the EMBL database under accession number AB011160. For the purpose of this application,

- 12 -

application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSSRP14A, unless stated otherwise or apparent from the context.

Marker 23 is bone small proteoglycan I (biglycan) protein. The gene for this protein has been cloned and sequenced (Fisher et al., J. Biol. Chem. 264:4571-4576 (1989)). These authors identify bone small proteoglycan I (biglycan) as a protein derived from human bone-derived cells. There is no teaching in this paper of an association with prostate cancer. The sequence of bone small proteoglycan I (biglycan) protein is disclosed in Fisher et al. and is present in the EMBL database under accession number HSHPGI. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSHPGI, unless stated otherwise or apparent from the context.

Marker 24 is the human KIAA0045 gene. The gene for this protein has been cloned and sequenced (Nomura N et al., DNA Res 1:223-229 (1994)). These authors identify KIAA0045 as a protein from the human immature myeloid cell line KG-1. There is no teaching in this paper of an association with prostate cancer. The sequence of KIAA0045 is disclosed in the EMBL database under accession number HSKG1C (accession no. D28476). For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSKG1C, unless stated otherwise or apparent from the context.

Marker 25 is b-cell receptor associated protein, also known as REA (repressor of estrogen activity). The gene for this protein has been cloned and sequenced (Montano et al., Proc Natl Acad Sci U.S.A. 96:6947-6952 (1999)). These authors identify REA as a protein derived from human breast. There is no teaching in this paper of an association with prostate cancer. The sequence of b-cell receptor associated /REA protein, is disclosed in Montano et al. and is present in the EMBL database under accession number AF150962. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number AF150962, unless stated otherwise or apparent from the context.

Marker 26 is the cystatin B protein. The gene for cystatin B has been cloned and sequenced (Pennacchio et al., Science 271:1731-1734 (1996)). These authors identify cystatin B protein as associated with progressive myoclonus epilepsy disease. There is no teaching in

- 14 -

markers such as PSA, prostatic acid phosphatase, prostate-specific membrane antigen or others (e.g. those reviewed in Gao et al., Prostate 31:264-281 (1977)); markers associated with cell proliferation, invasiveness, tumour angiogenesis, metastasis or any other aspect of carcinogenesis; or markers for other diseases and conditions. Levels of Marker mRNA in the test sample can be detected by any technique known in the art. These include Northern blot analysis, reverse transcriptase-PCR amplification (RT-PCR), microarray analysis and RNase protection.

In one embodiment, levels of Marker RNA in a sample can be measured in a Northern blot assay. Here, tissue RNA is fractionated by electrophoresis, fixed to a solid membrane support, such as nitrocellulose or nylon, and hybridised to a probe or probes capable of selectively hybridising with the Marker RNA to be detected. The actual levels may be quantitated by reference to one or more control housekeeping genes. Probes may be used singly or in combination. This may also provide information on the size of mRNA detected by the probe. Housekeeping genes are genes which are involved in the general metabolism or maintenance of the cell, and are considered to be expressed at a constant level irrespective of cell type, physiological state or stage in the cell cycle. Examples of suitable housekeeping genes are: beta actin, GAPDH, histone H3.3 or ribosomal protein L13 (Koehler et al., Quantitation of mRNA by Polymerase Chain Reaction. Springer-Verlag, Germany (1995)).

To gauge relative expression levels, a control sample can be run alongside the test sample or, the test result/value can be compared to Marker expression levels expected in a normal or control tissue. These control values can be generated from prior test experiments using normal or control tissues, to generate mean or normal range values for each Marker.

In another embodiment, the Marker nucleic acid in a tissue sample is amplified and quantitatively assayed. The polymerase chain reaction (PCR) procedure can be used to amplify specific nucleic acid sequences through a series of iterative steps including denaturation, annealing of oligonucleotide primers (designed according to the published Marker sequence to be detected), and extension of the primers with DNA polymerase (see, for example, Mullis, et al., U.S. patent No. 4,683,202; Loh et al., Science 243:217 (1988)). In reverse transcriptase-PCR (RT-PCR) this procedure is preceded by a reverse transcription step to allow a large amplification of the number of copies of mRNA (Koehler et al., *supra*).

Other known nucleic acid amplification procedures include transcription-based amplification systems (TAS) such as nucleic acid based sequence application (NASBA) and 3SR (Kwoh et

- 16 -

A variety of methods are currently available for making arrays of biological molecules. The 'dot or slot blot' approach, whereby an ordered array of DNA is vacuum blotted using a manifold, or hand blotted by capillary action, onto a porous membrane, such as nylon or nitrocellulose has been around for many years (Maniatis et al., Molecular Cloning-A Laboratory Manual, First Edition, Cold Spring Harbor, (1982)). Methods for preparing a plurality of oligonucleotide sequences and for attaching these to solid supports at high density are also known in the art. For example, US Patent No. 4,562,157 describes a method of using photo-activatable cross-linking groups to immobilise pre-synthesised ligands on surfaces. Fodor et al. (Nature 364:555-556 (1993)) and US Patent No. 5,143,854 describe the 'light-directed chemical synthesis' method for synthesising ligands, including oligonucleotides, directly onto a substrate surface at the desired location. US 5,700,637 also describes methods for *in situ* synthesis of oligonucleotides on solid support surfaces. In addition, such methods for preparing microarrays can easily be automated. International Publication No. WO 95/35505 discloses an automated capillary dispensing device and method for applying biological macromolecules to solid supports. International Publication No. WO 97/44134 also describes devices for delivery of small volumes of liquid (which may contain biological macromolecules) in a precise manner to produce micro-sized spots on a solid surface to generate a microarray. Similarly, International Publication No. WO 98/10858 also describes an apparatus for the automated synthesis of molecular arrays. Techniques exist for applying the oligonucleotides to the array at high density and for example, techniques exist for applying well in excess of 10^3 distinct polynucleotides per 1 cm^2 .

Microarray technology makes it possible to simultaneously study the expression of many thousands of genes in a single experiment. Analysis of gene expression in human tissue (e.g. biopsy tissue) can assist in the diagnosis and prognosis of disease and the evaluation of risk for disease. A comparison of levels of expression of various genes from patients with defined pathological disease conditions with normal patients enables an expression profile, characteristic of disease, to be created. There are currently two main approaches to analyse gene expression using microarrays. In the first approach, cDNA fragments, often generated by PCR, for each of the genes under study are attached to an array. Typically, mRNA isolated from the test samples (i.e. induced or un-induced) is reverse transcribed into cDNA with incorporation of a fluorescent label. The cDNA is sheared and hybridised to the array. If a control test sample is to be run at the same time, mRNA from this sample can be reverse

- 18 -

In transcript profiling, several or many mRNAs are detected in the same procedure. One or more of these mRNAs may be diagnostic of cancer cells (i.e prostate cancer) in a tissue sample. In one embodiment, combinations of probes can be used to classify the cancer cells into clinically relevant types, according to the complex expression pattern of Markers measured on the array. Such classification may help to define which tumours are growing aggressively, or harbour latent signs of aggression, or are less aggressive or benign. The array provides a quantitative measure of Marker RNAs. This is done by comparison of Marker RNA signal with control signal. In a preferred embodiment hybridisation signals generated are measured by computer software analysis of images on phosphorimage screens exposed to radioactively labelled tissue RNA hybridised to a microarray of probes on a solid support such as a nylon membrane. In another, quantities are measured by densitometry measurements of radiation-sensitive film (e.g. X-ray film), or estimated by visual means. In another embodiment quantities are measured by use of fluorescently labelled probe, which may be a mixture of tumour and normal RNA differentially labelled with different fluorophores, allowing quantities of Marker mRNAs to be expressed as a ratio versus the normal level. The solid support in this type of experiment is generally a glass microscope slide, and detection is by fluorescence microscopy and computer imaging.

The detection of specific interactions may be performed by detecting the positions where the labelled target sequences are attached to the array. Radiolabelled probes can be detected using conventional autoradiography techniques. Use of scanning autoradiography with a digitised scanner and suitable software for analysing the results is preferred. Where the label is a fluorescent label, the apparatus described, e.g. in International Publication No. WO 90/15070, US Patent No. 5, 143,854 or US Patent No. 5,744,305 may be advantageously applied. Indeed, most array formats use fluorescent readouts to detect labelled capture:target duplex formation. Laser confocal fluorescence microscopy is another technique routinely in use (M.J.Kozal et al., Nature Medicine 2:753-759 (1996)). Mass spectrometry may also be used to detect oligonucleotides bound to a DNA array (Little et al, Analytical Chemistry 69: 4540-4546, (1997)). Whatever the reporter system used, sophisticated gadgetry and software may be required in order to interpret large numbers of readouts into meaningful data (such as described, for example, in US Patent No. 5,800,992 or International Publication No. WO 90/04652).

- 20 -

RNA can be extracted at a later date. Proprietary reagents are available which allow tissue or cells to be conveniently stored for several days at room temperature and up to several months at 4°C (e.g. RNAlater, Ambion Inc., TX). Prior to extraction, methods such as grinding, blending or homogenisation are used to dissipate the tissue in a suitable extraction buffer.

- 5 Typical protocols then use solvent extraction and selective precipitation techniques. Example 1 describes such a method.

In another embodiment, tissue is directly analysed for the presence of Marker nucleic acid. This can be by *in situ* hybridisation, where sections of tissue may be interrogated with specific probes to determine which morphological cell type in the sample displays a marker

10 nucleic acid, such as sequences corresponding to Markers 1-26. *In situ* hybridisation typically comprises 3 steps. Firstly tissue is fixed, and sections are prepared by standard treatments known to those in the art (Polack and McGee, *In situ* hybridisation: principles and practice. Oxford University Press, 1998). Secondly, Marker mRNA and amplified Marker DNA can be detected by hybridisation with e.g. a biotin-, digoxigenin- or radio-labelled Marker probe,

15 typically for 2-16 hours at 42°C in a suitable hybridisation buffer. A typical buffer might contain 50% formamide, 5% dextran sulphate, 2xSSC and 10-20ng of probe per 7µl (Herrington and McGee, *Diagnostic Molecular Pathology*. IRL Press, Oxford, 1992). The probe can be made of DNA or RNA. Lastly, following stringency washes, the probes in hybridisation complexes are detected with chromogenic or fluorescent reagents, which can be

20 visualised by microscopy, or by autoradiography in the case of radiolabelled probes. Signal amplification systems using e.g. tyramide can be used to increase sensitivity (Polack and McGee, *supra*). *In situ* PCR using oligonucleotide probes complementary to the nucleic acid of any of Markers 1 - 26 is therefore envisaged. *In situ* hybridisation can follow *in situ* PCR, giving greater specificity (Polack and McGee, *supra*). Techniques for quantitation of signal,

25 and quantification of positive cells in a section are available to the pathologist using image analysis. *In situ* Marker visualisation permits localisation of signal in mixed-tissue specimens commonly found in tumours, and is compatible with many histological staining procedures.

In one embodiment, several probes can be differentially labelled and hybridised simultaneously to the same section, and detected using appropriate reagents. In another, serial

30 sections from the same sample can be analysed with a panel of probes. Quantitation may involve comparison with one or more control housekeeping genes as discussed above.

- 22 -

cDNA) then the probe or primer sequence can hybridise to the sense or antisense strand. If however the target is mRNA (single stranded sense strand) the primer/probe sequence will have to be the antisense complement.

5 An example of a suitable hybridisation solution when a nucleic acid is immobilised on a nylon membrane and the probe nucleic acid is greater than 500 bases or base pairs is: 6 x SSC (saline sodium citrate), 0.5% SDS (sodium dodecyl sulphate), 100µg/ml denatured, sonicated salmon sperm DNA. The hybridisation being performed at 68°C for at least 1 hour and the filters then washed at 68°C in 1 x SSC, or for higher stringency, 0.1 x SSC/0.1% SDS.

10 An example of a suitable hybridisation solution when a nucleic acid is immobilised on a nylon membrane and the probe is an oligonucleotide of between 12 and 50 bases is: 3M trimethylammonium chloride (TMACl), 0.01M sodium phosphate (pH 6.8), 1mM EDTA (pH 7.6), 0.5% SDS, 100µg/ml denatured, sonicated salmon sperm DNA and 0.1 dried skimmed milk. The optimal hybridisation temperature (T_m) is usually chosen to be 5°C below the T_i of
15 the hybrid chain. T_i is the irreversible melting temperature of the hybrid formed between the probe and its target. If there are any mismatches between the probe and the target, the T_m will be lower. As a general guide, the recommended hybridisation temperature for 17-mers in 3M TMACl is 48-50°C; for 19-mers, it is 55-57°C; and for 20-mers, it is 58-66°C.

A suitable hybridisation protocol is described in Example 5, however, operable
20 variations to this method will be apparent to the person skilled in the art.

Where the cDNA molecules of Markers 1 - 26 encode proteins or parts of cellular proteins, these may themselves act as prostate disease Markers. To detect proteins in tissue, cells, body fluids or extracts of these sample types, specific antibody can be used. These antibodies can be prepared using the Marker protein/polypeptides.

25 Methods of making and detecting labelled antibodies are well known (Campbell; Monoclonal Antibody Technology, in: Laboratory Techniques in Biochemistry and Molecular Biology, Volume 13. Eds: Burdon R et al. Elsevier, Amsterdam (1984)). The term antibody includes both monoclonal antibodies, which are a substantially homogeneous population, and polyclonal antibodies which are heterogeneous populations. The term also includes inter alia,
30 humanised and chimeric antibodies. Monoclonal antibodies to specific antigens may be obtained by methods known to those skilled in the art, such as from hybridoma cells, phage display libraries or other methods. Monoclonal antibodies may be inter alia, human, rat or

- 24 -

that the nucleic acid sequence encoding the polypeptide is transcribed into RNA in the host cell transformed or transfected by the expression vector construct. The coding sequence may or may not contain a signal peptide or leader sequence for secretion of the polypeptide out of the host cell. Expression and purification of the Marker polypeptides can be easily performed using methods well known in the art (for example as described in Sambrook et al. *supra*).

The Marker polypeptides so produced can then be used to inoculate animals, from which serum samples, containing the specific antibody against the introduced Marker protein/polypeptide, can later be obtained.

Rodent antibodies may be humanised using recombinant DNA technology according to techniques known in the art. Alternatively, chimeric antibodies, single chain antibodies, Fab fragments may also be developed against the polypeptides of the invention (Huse et al., Science 256:1275-1281 (1989)), using skills known in the art. Antibodies so produced have a number of uses which will be evident to the molecular biologist or immunologist skilled in the art. Such uses include, but are not limited to, monitoring enzyme expression, development of assays to measure enzyme activity and use as a therapeutic agent. Enzyme linked immunosorbant assays (ELISAs) are well known in the art and would be particularly suitable for detecting the Marker proteins or polypeptide fragments thereof in a test sample.

The expression system described in Example 3 can produce protein for use as an antigen for the generation of antibodies for use in an ELISA assay to detect Marker protein in body fluids or by immunohistochemistry (as described in Example 4) or other means. In addition, an antibody could be used individually or as part of a panel of antibodies, together with a control antibody which reacts to a common protein, on a dipstick or similar diagnostic device.

Levels of Marker gene expression can also be detected by screening for levels of polypeptide (Marker protein). For example, monoclonal antibodies immunoreactive with a Marker protein can be used to screen a test sample. Such immunological assays can be done in any convenient format known in the art. These include Western blots, immunohistochemical assays and ELISA assays. Functional assays can also be used, such as protein binding determinations.

In another preferred embodiment antibodies directed against a Marker protein or proteins can be used, to detect, prognose, diagnose and stage prostate cancer or its precursor lesions, or related prostate disorders. Various histological staining methods known in the art,

- 26 -

Thus, according to another aspect of the invention there is provided a method for treating a patient suffering from a prostate disorder comprising administering to said patient an effective amount of an anti-sense molecule capable of binding to the mRNA of a Marker gene selected from the group consisting of: Marker 1 - Marker 26, and inhibiting expression
5 of the protein product of the Marker gene.

Complete inhibition of protein production is not essential, indeed may be detrimental. It is likely that inhibition to a state similar to that in normal tissues would be desired.

This aspect of antisense therapy is particularly applicable if the prostate disorder is a direct cause of over-expression of the Marker gene(s) in question, although it is equally
10 applicable if said Marker gene(s) indirectly cause the prostate disorder. Having identified the particular Marker genes (1 - 26) over-expressed in prostate disorders, and with knowledge of the gene and mRNA sequence the person skilled in the art is able to design suitable antisense nucleic acid therapeutic molecules and administer them as required.

Antisense oligonucleotide molecules with therapeutic potential can be determined
15 experimentally using well established techniques. To enable methods of down-regulating expression of a Marker gene of the present invention in mammalian cells, an example antisense expression construct can be readily constructed for instance using the pREP10 vector (Invitrogen Corporation). Transcripts are expected to inhibit translation of the gene in cells transfected with this type of construct. Antisense transcripts are effective for inhibiting
20 translation of the native gene transcript, and capable of inducing the effects (e.g., regulation of tissue physiology) herein described. Oligonucleotides which are complementary to and hybridisable with any portion of Marker gene mRNA are contemplated for therapeutic use. U.S. Patent No. 5,639,595, "Identification of Novel Drugs and Reagents", issued Jun. 17, 1997, wherein methods of identifying oligonucleotide sequences that display *in vivo* activity
25 are thoroughly described, is herein incorporated by reference. Expression vectors containing random oligonucleotide sequences derived from Marker polynucleotides are transformed into cells. The cells are then assayed for a phenotype resulting from the desired activity of the oligonucleotide. Once cells with the desired phenotype have been identified, the sequence of the oligonucleotide having the desired activity can be identified. Identification may be
30 accomplished by recovering the vector or by polymerase chain reaction (PCR) amplification and sequencing the region containing the inserted nucleic acid material. Antisense molecules can be synthesised for antisense therapy. These antisense molecules may be DNA, stable

- 28 -

analogous to quantitative RT-PCR already described. Marker gene amplification could also be detected *in situ* in cells by probe hybridisation. Fluorescently-labelled probes (used in Fluorescent in situ Hybridisation or FISH) are ideal for this purpose as they allow subnuclear localisation via microscopy, and permit estimates of gene copy number by subnuclear spot intensity or number (Mark et al. Exp Mol Pathol 66:170-8 (1999)). Quantitative analysis of DNA for amplifications of Marker genes can also be carried out by Southern analysis, a method which is widely known to those skilled in the art (Sambrook et al., *supra*). DNA can be extracted from clinical material using established methods (Sambrook et al., *supra*). The methods of the invention can therefore also be directed to measuring genomic DNA levels of one or more of the identified Markers.

If a Marker mRNA encodes a secreted protein, that protein is likely to be present in body fluids. Proteins, especially secreted proteins, can be quantified in blood, serum, urine, semen and other fluids. Specific antibodies can often detect abundant proteins in ELISA tests on body fluid samples without enrichment. Prostate-specific antigen (PSA) falls into this category. PSA is an important Marker produced by prostatic epithelial cells and almost always expressed in prostate cancer, though not exclusively so. Clinically significant levels of proteins such as PSA are defined by appropriate studies, and protein levels are typically given as ng per mL of sample. Serum protein tests with greater specificity for cancer are needed. Detection of rare proteins may require that the protein is concentrated by e.g. precipitation. Thus, in a further embodiment of the invention diagnosis or prognosis or prostate disorder, or stage monitoring of the prostate disorder or therapeutic efficacy assessment is performed by testing for aberrant levels of one or more Marker proteins of the invention, which Marker protein is a secreted protein, in a bodily fluid.

The inventors predict that Markers 9, 19 and 24 (at least) are secreted proteins.

All the essential materials and reagents required for detecting or monitoring prostate disorder Markers in a test sample may be assembled together in a kit. Such a kit may comprise one or more diagnostic cDNA probes or oligonucleotide primers together with control probes/primers. The kit may contain probes immobilised on a microarray substrate such as a filter membrane or silicon-based substrate. The kit may also comprise samples of total RNA derived from tissues of various physiological states, such as normal, BPH, confined tumour and metastatic tumour, for example, to be used as controls. The kit may also comprise appropriate packaging and instructions for use in the methods of the invention.

- 30 -

Elmer 7700). Preferred oligonucleotide primers for detection of Marker mRNAs are selected from Markers 1-26.

In another embodiment the kit comprises one or more antibodies specific for one or other of the Markers identified herein for use in immunohistochemical analysis.

5 In another embodiment the kit is an ELISA kit comprising one or more antibodies specific for one or other of the Markers identified herein.

In another aspect of the invention, one or more of the 26 Markers can also be used in biochemical assays to identify agents which modulate the activity of the Marker proteins. The design and implementation of such assays will be evident to the biochemist of ordinary skill.

10 The protein, particularly if it is a biochemical enzyme, may be used to turn over a convenient substrate whilst incorporating or losing a labelled component to define a test system. Test compounds are introduced into the test system and measurements made to determine their effect on enzyme activity. Such assays are useful to identify inhibitors of the enzyme which may prove valuable as therapeutic agents.

15 The inventors believe that Marker 19 (Factor V) is an example of a suitable biochemical enzyme that can be used in a suitable biochemical assay to identify modulators.

In a further aspect of the invention, each of the Markers can be used to characterise cell cultures in a screen for therapeutic agents, such as a high throughput screen. Effects of test compounds may be assayed by changes in mRNA or protein of any of Markers 1-26. As
20 described above, cells (i.e. mammalian, bacterial etc) can be engineered to express one of the Markers identified herein.

Thus, according to a further aspect of the invention there is provided a method of testing potential therapeutic agents for the ability to suppress a prostate disorder phenotype comprising contacting a test compound with a cell engineered to express one of the Markers
25 identified herein; and determining whether said test compound suppressed expression of the Marker.

Thus, according to a further aspect of the invention there is provided a screening assay or method for identifying potential anti-prostate disorder therapeutic compounds comprising contacting an assay system capable of detecting the effect of a test compound against
30 expression levels of one or more of the Markers selected from the group consisting of: Marker 1 - Marker 26, with a test compound and assessing the change in expression level of the particular Marker under study.

- 32 -

Further features of the invention include:

A method of treatment of a patient suffering from a prostate disorder, comprising administration to the patient of a compound capable of reducing the transcription or expression of any one of Markers 1 - 26.

- 5 A method of treatment of a patient suffering from a prostate disorder, comprising administration to the patient an antisense nucleic acid molecule targeted against the mRNA of any one of Markers 1 - 26.

Use of an antisense nucleic acid molecule or an antibody directed against any one of Markers 1 - 26, in the manufacture of a medicament for treating a prostate disorder.

- 10 Each aspect of the invention involves detection or use of one or more of Markers 1 - 26. A preferred sub-group of Markers are those selected from the group consisting of Markers: 8, 13, 14, 15, 17, 18, 19, 20, 22, 24, 25 and 26.

The invention will be further described by way of the following non-limiting examples and figures in which data illustrating over-expression of markers is included:

- 15 **Figure 1.** Filter microarray data showing over-expression of Markers 1 - 18 and 20-25 in prostate cancer relative to normal prostate. Overexpression value for Marker 26 (not shown) was found to be at the same level as Marker 13. Values given are mean expression level for 9 prostate cancer samples in those samples where over-expression of 1.5-fold or more was detected. Expression is given relative to normal prostate level (i.e. compared to the mean of 3
20 normal prostate datasets. These 3 datasets comprised 2 different RNA mixtures, each with at least 10 normal sample components).

Figure 2. Filter microarray data showing over-expression of Markers 1-5, 7-12, 17, 18, 20 and 23 in prostate cancer relative to BPH. Values given are mean expression level for 9 prostate cancer samples in those samples where over-expression of 1.5-fold or more was
25 detected. Expression is given relative to BPH level (i.e. compared to the mean of 13 BPH datasets).

Example 1: Identification and Evaluation of Markers of Prostate Disease by cDNA

Microarray Analysis

- 30 The microarray analysis protocol described in the following example was developed as a means to determine the relative abundances of mRNA species that are expressed in various tissues. Microarray analysis was used to identify differentially expressed RNA species

- 34 -

treated with DNase I (also Gibco BRL), again as recommended by the manufacturer. For 1st strand label synthesis, 20µg total RNA was reverse transcribed into cDNA in the presence of radiolabelled nucleotide (³³P), using Superscript II enzyme (Gibco BRL) according to manufacturer's instructions. Labelled RNA was purified using GFX columns (Amersham Pharmacia Biotech, St. Albans, UK) according to manufacturers instructions, and added to 10 ml/filter of Church hybridisation buffer described below.

Prior to hybridisation, filters were wetted briefly in 2xSSC, then incubated in 5ml prewarmed Church hybridisation solution (0.5M sodium phosphate, 7% SDS, 1mM EDTA, pH 7.2) at 65°C for 2-6 hours. Probe was denatured at 100°C for 5 min, placed on ice for at least 5 min, then added to 5ml fresh Church hybridisation buffer in the presence of the filter, and mixed by swirling the bottle with the lid on for even probe distribution. Hybridisation was carried out at 65°C for 12-16 hours. Filters were rinsed briefly in pre-warmed Church wash solution (40mM sodium phosphate, 1% SDS, pH 7.2), then incubated twice in the same solution at 65°C for 20 min. Filters were drained briefly, wrapped in Saran wrap (Dow Chemical Company, USA) and exposed to phosphor screens for 3-6 days (screens and cassettes supplied by Molecular Dynamics, CA). The screens were then scanned using a Storm 830 phosphorimager (Molecular Dynamics). Array Vision software (Incyte Pharmaceuticals, CA, USA) was used to visualise the hybridisation images and generate quantitative numbers for each spot.

Typically, for comparison of data from different arrays, data is generated as a value relative to an internal standard for each array. Following export of ArrayVision data into Microsoft Excel format, spot measurements for data were normalised to housekeeping gene hybridisation signals known to be constant or relatively constant, using a simple Excel macro. A total of 38 spot values (from 16 different clones) representing 12 different housekeeping genes were used to generate an average housekeeping measurement. The gene and EMBL accession number of the housekeeping genes used are listed in Table 1. Background values were not subtracted, but to eliminate spurious low-level signals indistinguishable from noise, a local background value from 3 blank spots out of every 7x7 array was taken, and every value <2-fold this value was ignored.

To generate representative values for normal prostate, data from 3 array hybridisation experiments to normal prostate RNA was normalised to the mean housekeeping gene value, then the mean of these 3 values was taken as the control dataset.

- 36 -

histopathological analysis. Table 2 lists the 26 genes we have identified as being Markers for prostatic disease.

Control number	Gene	NCBI genInfo identifier	Spots on Array
1	Human ribosomal protein S14 gene	g337498	2
2	Human ribosomal protein S14 gene	g337498	2
3	Human acidic ribosomal phosphoprotein P0	g190231	2
4	Human alpha-tubulin mRNA	g340020	2
5	Human mRNA for TEF-5 protein	g1848081	2
6	Human transcriptional enhancer factor	g339440	2
7	Human transcription factor RTEF-1	g1561727	2
8	Human hypoxanthine phosphoribosyltransferase	g184349	2
9	Human hypoxanthine phosphoribosyltransferase	g184349	2
10	Human mRNA for transcription factor, TEF3	g1403337	2
11	Human acidic ribosomal phosphoprotein P1	g190233	2
12	Human mRNA for ribosomal protein L19	g36127	4
13	Human hH3.3B gene for histone H3.3	g761715	4
14	Human pancreatic phospholipase A-2 (PLA- 2)	g190008	2
15	Human mRNA for ribosomal protein L19	g36127	4
16	Human acidic ribosomal phosphoprotein P1	g190233	2

5 Table 1. Names and identities of control housekeeping genes used to normalise microarray data. This set was identified from a wider set, by comparison of variability of candidate housekeeping gene signals over 13 filter hybridisation experiments as a group showing minimal variability.

- 38 -

prostate cancer compared to BPH, or in the case of Marker 19, not detected at all in normal prostate. In addition, Markers 6, 13, 14 and 25 were not detected in BPH.

Example 2: Quantitative RT-PCR test

5 RT-PCR is a convenient method for assaying the relative abundance of mRNA species expressed in various cells, tissues and organs.

Reproducibly accurate amplification of an mRNA can be achieved by firstly synthesising a DNA template from the mRNA using a reverse transcriptase, then by PCR using a heat-stable DNA polymerase (e.g. Taq polymerase). An optimised amplification
10 protocol is essential for quantitation of nucleic acids by PCR because small differences in efficiency of the reaction can greatly influence the reaction rates, with a subsequent effect on PCR yield. Optimisation of PCR is not usually problematic (refer to: Koehler *supra*, and references therein). Quantitation of RT-PCR products can be done while the reaction products are building up exponentially, preferably following each round of amplification
15 (known as real-time PCR). For quantitation, analysis is carried out by reference to one or more housekeeping genes which are also amplified by RT-PCR, often in a different reaction tube. Quantitation of RT-PCR product may be undertaken, for example, by gel electrophoresis visual inspection or image analysis, HPLC (Koehler et al., *supra*) or by use of suitable detection methods such as described above.

20 In practise, specific pairs of oligonucleotide primers corresponding to Markers 1-26 can be used to trigger amplification of specific marker mRNAs. The Marker mRNA is quantitated relative to a standard housekeeping mRNA using a real-time RT-PCR assay. For the purpose of this example, the housekeeping gene is ribosomal protein L19, but any such gene could be used.

25 Firstly, RNA is isolated from a clinical or tissue sample as in example 1. Aliquots of RNA are used in a reverse transcription reaction using random hexamers or oligo-dT as primers, and for example Moloney Murine Leukaemia Virus (MuLV) or Avian Myeloblastosis Virus (AMV) reverse transcriptase. A typical 20ml reverse transcription reaction comprises the following: 10 mM each dNTP, 50 mM Tris-HCl, pH8.3, 50 mM KCl,
30 50 mM dithiothreitol, 0.5 mM spermidine, 10ng oligo-dT primer, up to 1µg RNA, dissolved in DEPC-treated water. This mixture is incubated at 65°C for 15 min, then chilled on ice for 5 min. 5 units of AMV reverse transcriptase and 10 units of RNase inhibitor are added. The

- 40 -

Example 3: Generation of antibody specific to Protein encoded by Markers 1-26

Antibody specific to Markers 1-26 could be used to detect protein products of these Markers in tissue samples and body fluids. Ideally for diagnosis, easily accessible samples are required, involving minimally invasive techniques. Typically, these samples include blood, plasma, serum, saliva, urine and semen. These fluids are far removed from prostate cells and, in a normal individual, would not be expected to contain mRNA of prostatic origin. However, protein may be transported across cell membrane barriers into body fluids. Prostate-specific antigen is one such protein, and is easily assayed by a simple blood test using specific antibody against the protein. For new markers, e.g. Markers 1-26, it may be necessary to generate novel antibody. This example describes a method of generating protein in vitro, which can then be used to raise antibody.

One way to generate protein is to express the cloned gene in *E. coli*. This requires that the cDNA, e.g. Markers 1-26 is cloned into a vector capable of expressing the encoded protein in the bacterial host. Optionally, it may be desirable to incorporate a molecular tag into the protein so that it can be easily purified. One such tag in wide use is the 6xHis tag. A protein with this tag is easily isolated from a cell culture extract by affinity chromatography, and can be used in relatively pure form to inoculate an animal for antibody generation.

Having subcloned the cDNA or cDNA fragment into an appropriate vector, e.g. pHAT10, 11, 12 or 20 (Clontech, Palo Alto, CA) and confirmed the integrity of the insert such as sequence and orientation. The subclone is cultured on a suitable scale, cells are harvested by centrifugation and then lysed, and the protein extract incubated with TALON resin according to manufacturer's instructions (Clontech, Palo Alto, CA). The resin is washed and recombinant protein is eluted by adjusting the pH or imidazole concentration. If desired, the 6xHIS tag can be enzymatically removed using a specific protease (Clontech). To generate a specific antibody, the purified protein is injected into a host animal, usually rabbit, sheep or goat. After boost injections, the serum is periodically collected and tested for antibody.

Polyclonal antibody can be purified from the serum using standard techniques, or the animal's spleen can be harvested for the production of hybridomas. Techniques for the production of antibodies, both polyclonal and monoclonal, are well known to those skilled in the art (Catty D (ed.) Antibodies: A Practical Approach. Vol 1 (1988), Vol2 (1989)).

- 42 -

A hybridisation probe can be generated from a synthetic oligonucleotide or a dephosphorylated restriction fragment sequence by addition of a radioactive 5' phosphate group from [γ - ^{32}P]ATP by the action of T4 polynucleotide kinase. 20 pmoles of the oligonucleotide are added to a 20 μl reaction containing 100mM Tris, pH7.5, 10mM MgCl_2 , 0.1mM spermidine, 20mM dithiothreitol (DTT), 7.55 μM ATP, 55 μCi [γ - ^{32}P]ATP and 2.5u T4 polynucleotide kinase (Pharmacia Biotechnology Ltd, Uppsala, Sweden). The reaction is incubated for 30 minutes at 37°C and then for 10 minutes at 70°C prior to use in hybridisation. Methods for the generation of hybridisation probes from oligonucleotides or from DNA and RNA fragments are described in Chapters 11 and 10 respectively in Sambrook et al. (*ibid*). A number of proprietary kits are also available for these procedures.

Filter preparation

The sample DNA could be isolated and run on an agarose gel and Southern blotted onto a nitrocellulose or nylon filter using standard techniques.

Hybridisation conditions

Filters containing the nucleic acid are pre-hybridised in 100ml of a solution containing 6x SSC, 0.1%SDS and 0.25% dried skimmed milk (Marvel™) at 65°C for a minimum of 1 hour in a suitable enclosed vessel. A proprietary hybridisation apparatus such as model HB-1 (Techne Ltd) provides reproducible conditions for the experiment.

The pre-hybridisation solution is then replaced by 10ml of a probe solution containing 6xSSC, 0.1% SDS, 0.25% dried skimmed milk (e.g. Marvel™) and the oligonucleotide probe generated above. The filters are incubated in this solution for 5 minutes at 65°C before allowing the temperature to fall gradually to below 30°C. The probe solution is then discarded and the filters washed in 100ml 6xSSC, 0.1% SDS at room temperature for 5 minutes. Further washes (1-3) are then made in fresh batches of the same solution at 30°C and then, optionally, in 10°C increments up to 60°C for 5 minutes per wash.

After washing, the filters are dried and used to expose an X-ray film such as Hyperfilm™ MP (Amersham International) at -70°C in a light-tight film cassette using a fast tungstate intensifying screen to enhance the photographic image. The film is exposed for a suitable period (normally overnight) before developing to reveal the photographic image of the radio-active areas on the filters. Related nucleic acid sequences are identified by the presence of a photographic image compared to totally unrelated sequences which should not

- 44 -

then chilled quickly on ice-water, is placed on the tissue section and a coverslip is sealed on to the slide using rubber solution. A suitable probe is a digoxigenin-labelled 300-600 base pair cDNA clone or fragment thereof, corresponding to one of Markers 1-26, prepared by one of several methods known to the art. The slides are then incubated in a sealed chamber for 16 h at 37°C. The rubber solution is peeled away and coverslips are removed by immersion of the slides in 2x SSC, 0.1% SDS. Slides are washed 4 times for 5 min each at room temperature, and twice for 10 min at 65°C, in 2x SSC, 0.1% SDS, and rinsed briefly in 2x SSC. Hybridisation is detected as follows: Nonspecific binding is blocked by incubating the slides in blocking buffer (0.1M Tris-HCL pH 7.5, 0.1M NaCl, 2mM Mg Cl₂, 3% bovine serum albumin) for 10 min. Slides are then flooded with anti-digoxigenin-conjugated alkaline phosphatase (1/500 dilution of stock; Boehringer, Mannheim, Germany) in blocking buffer, and incubated 2 h at room temperature, then washed in blocking buffer 3 times for 3 min each. Slides are placed in buffer 2 (0.1 M Tris-HCl pH 9.5, 0.1 M NaCl, 50 mM MgCl₂) for 10 min, then immersed in substrate buffer (100mL buffer 2 containing 25 mg levamisole, plus 35 mg nitroblue tetrazolium chloride dissolved in 277 mL 70% dimethylformamide, plus 17 mg 5-bromo-4-indoyl-phosphate dissolved in 222 mL 100% dimethylformamide) for 10-30 min in the dark. The slides are then immersed in 20 mM Tris-HCl pH7.5, 5 mM EDTA for 5 min, rinsed with tap water for 5-10 min, mounted in aqueous mountant and examined under a microscope.



Example 7. Screening for therapeutic agents capable of inhibiting expression of any of Markers 1-26.

Markers 1-26 may be employed in a process for screening compounds which either inhibit, promote or modulate the expression of Markers 1-26. Examples of potential Marker 1-26 agonists are small molecules such as organic molecules or peptides, antibodies or oligonucleotides which bind to Markers 1-26 and inhibit expression or activity. One assay for therapeutic agents uses cultured cells, and measures transcript abundance using a microarray as described in Example 1. Typical prostate cell lines such as LNCaP, PC-3 and DU145 are available from the European Collection of Cell Cultures, Salisbury UK. Cell culture is a standard technique well known to those practised in the art. Briefly, in one suitable example, a seed vial of LNCaP cells is inoculated into 500mL culture flasks containing 20-50ml RPMI medium containing 10% fetal calf serum and 1% glutamate (all components Gibco, UK).

Claims:

1. A method for diagnosing or prognosing or monitoring a prostate disorder comprising testing a biological sample from an individual for aberrant levels of one or more of the
5 Markers selected from the group consisting of: Marker 1 to Marker 26.
2. A method as claimed in claim 1, wherein the Marker(s) are selected from the group consisting of Markers: 8, 13, 14, 15, 17, 18, 19, 20, 22, 24, 25 and 26.
- 10 3. A method for distinguishing prostate cancer from BPH comprising testing a biological sample for aberrant levels of one or more of the Markers selected from the group consisting of Marker 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 17, 18, 20, 23, 25 and 26.
4. A method as claimed in any of claims 1-3 wherein mRNA transcript levels and/or
15 protein levels of the Marker(s) is/are measured.
5. A method as claimed in claim 4, wherein mRNA transcript levels are measured using reverse-transcriptase polymerase chain reaction (RT-PCR).
- 20 6. A method for measuring Marker mRNA transcript levels as claimed in claim 4, which method involves the use of one or more oligonucleotide probes each capable of selectively hybridising to nucleic acid of a Marker of interest to determine the expression level of said Marker of interest.
- 25 7. A method for measuring Marker protein levels as claimed in claim 4, which method involves the use of one or more antibodies each capable of selectively binding to a Marker protein or protein fragment of interest to determine the expression level of said Marker of interest.
- 30 8. Use of an antibody selective for a Marker protein selected from the group consisting of: Marker 1 - Marker 26, in an assay to monitor therapeutic efficacy.

9. A diagnostic kit for diagnosing or prognosing or monitoring a prostate disorder comprising, one or more diagnostic probe(s) and/or one or more diagnostic primer(s) and/or one or more antibodies capable of selectively hybridising or binding to one or more of the Markers 1 -26.
- 5
10. A screening assay for identifying potential anti-prostate disorder therapeutic compounds comprising contacting an assay system capable of detecting the effect of a test compound on expression levels of one or more of the Markers selected from the group consisting of: Marker 1 - Marker 26, with a test compound and assessing the change in
10 expression level of the particular Marker under study.
11. A method of testing potential therapeutic agents for the ability to suppress a prostate disorder phenotype comprising contacting a test compound with a cell engineered to express one of the Markers identified herein; and determining whether said test compound suppressed
15 expression of the Marker.
12. A compound or agent identified by the screening assay according to claim 10 or the method according to claim 11.
- 20 13. A method of treatment of a patient suffering from a prostate disorder, comprising administration to the patient of a compound capable of reducing the transcription or expression of any one of Markers 1 - 26.

Figure 1**Figure 2**

- 2 -

5 cctccaccct ggtccgctc cctcgctgg ctcaaccgag tgcctccgac cccctcctc 840
agccctcccc caccacacag cccagcctcc tcggtctcct gtctcgttgc tgcttctgcc 900
tgtgtgtgg gggagagagg ccgcagccag gcctctgctg cccttctctgt gccccccagg 960
ttctatctcc ccgtcacacc cgaggcctgg cttcaggagg gagcggagca gccattctcc 1020
aggccccgtg gttgccccg gacgtgtgcy tctgctgctc cggggtgag ctggggtgtg 1080
ggatgcacgg cctcgtgggg gccgggcccgt cctccagccc cgtgctccc tggccagccc 1140
ccttgtcgtg gtcgggtccc tctaaccatg atgccttaac atgtggagtg taccgtgggg 1200
cctcactagc cctactccc tgtgtctgca tgagcatgtg gcctccccgt cccttccccg 1260
gtggcgaacc cagtgaacca gggacacgtg ggggtgtgctg gtgctgctcc ccagcccacc 1320
10 aatgcctggc cagcctgccc ccttccctgg acagggctgt ggagatggct ccggcggctt 1380
ggggaaagcg aaattgcaa cactcaagtc acctcagtag catccaggag gctgggtatt 1440
gtctgcctc tgccttttct gtctcagcgg cagtgccag agcccacacc ccccaagag 1500
ccctcgatgg acaggcctga cccaccccac ctggggccag ccaggagccc cgcctgggcc 1560
atcagtattt attgcctccg tccgtgccgt cctggggcca ctggctggcg cctcttcccc 1620
15 cagcctctca gtgccaccac ccccgccagc cttccctgac ccagccagga caaacaaggg 1680
accaagtga cacattgctg agagccgtct cctataggtc ccccgcccca tccccggtgt 1740
tgggtgtgtg tctgccaggc tcaggcagag gcgcctgtcc ctgcttcttt tctgaccggg 1800
aaataaatgc ccctgaagg 1819

20

<210> 3
<211> 996
<212> DNA
<213> Homo sapiens

25

<400> 3
agcttctctg atccctagta taaacacttc agtgttcccc tttcagttctt actactttga 60
ccgcgatgat gtggctttga agaactttgc caaatacttt cttcaccaat ctcatgagga 120
gagggaacat gctgagaaac tgatgaagct gcagaaccaa cgagggtggc gaattcttct 180
30 tcaggatata aaggtgaaca aaagatccta ggggtgtcat acttcatcat ctggcagtg 240
tcgggtatca gaaatcactt aaactagcaa ttgcccttat aaagtgatga tacactgggc 300
ttttgccttt tgtgcttttt taggcttacc atctaaacta aattaggcaa atagtaattg 360
cccttttgcc aaaacgtggg ggttagagat gatgggcttg ctgacttcta ggtagttgg 420
tagagatgca ttaacctatt ctcatcaga aaccagactg tgatgactgg gagagcgggc 480
35 tgaatgcaat ggagtgtgca ttacatttgg aaaaaaatgt gaatcagtca ctactggaac 540
tgcacaaact ggccactgac aaaaatgacc ccatgtgag tattggaacc ccaggaaata 600
aatggaggaa atcatttgcc ttagggtatg ggaagctgc ccactaactg tcttgcacca 660
ttgttttgca gttgtgtgac ttcattgaga cacattacct gaatgagcag gtgaaagcca 720
tcaaagaatt gggtgaccac gtgaccaact tgcgcaagat gggagcggcc gaatttggt 780
40 tggcgaata tctctttgac aagcacaccc tgggagacag tgataatgaa agctaagcct 840
cgggctaatt tccccatagc cgtggggtga cttccctggt caccaaggca gtgcatgcat 900
gttgggggtt cctttacctt ttctataagt tgtacaaaa catccactta agttctttga 960
ttgtaccat tccttcaaat aaagaaattt ggtacc 996

45

<210> 4
<211> 2353
<212> DNA
<213> Homo sapiens

50

<400> 4
gggttgagca caaggaggag aacgaccaca aagtcttcta cgggggtgac ctgaaagtgg 60
actgtgtggc caccgggctt cccaatcccc agatctcctg gagcctccca gacgggagtc 120
tggtgaactc cttcatgcag tcggatgaca gcgggtggag caccaagcgc tatgtcgtct 180
55 tcaacaatgg gacactctac tttaacgaag tggggtatgag ggaggaagga gactacacct 240
gctttgtctga aaatcaggtc gggaaaggag agatgagagt cagagtcaag gtggtgacag 300
cgccccccac catccggaac aagacttgc tggcggttca ggtgccctat ggagacgtgg 360
tcaactgtagc ctgtgaggcc aaaggagaac ccatgcccac ggtgacttgg ttgtcccca 420
ccaacaaggat gatccccacc tcctctgaga agtatcagat ataccaagat ggcactctcc 480
60 ttattcagaa agccagcgt tctgacagcg gcaactacac ctgcttggtc aggaacagcg 540
cgggagagga taggaagacg gtgtggattc acgtcaacgt ccagccaccc aagatcaacg 600
gtaaccccaa ccccatcacc actgtgcggg agatagcagc cgggggcagt cggaaactga 660
ttgactgcaa agctgaagcc atccccaccc cgagggtgtt atgggctttt cccgagggtg 720
tggttctgcc agctccatac tatgaaacc ggatcactgt ccatggcaac ggttccctgg 780

<211> 5086

<212> DNA

<213> Homo sapiens

5 <400> 6
gcctgcgcgc cgcagcccgc ctcgggcggg aggcggggagg cgggaggccg ggcccaggcc 60
ggggcagccc ctccaccgca ccgtccctggg ccggtgcccc ggtccgagtc gccttccgcc 120
tgcccccccc gccaatcccc cgcgcgggca gccccagcca ggtcccgccg ccaggccggc 180
tcccccccc gctccgcccc cggagccgca gccccggccg ccacgcgcgt cgcctatgtg 240
10 tggctccgc agccggcgct ggggacgcgc gcggccgaga ctctggcctg cagtcgcgcg 300
cgccgcgcgc aggtgggtgt tggactctag accatgtgcc taggtagaag ttttccctt 360
ctccgcagct ctgtccccct agcaacgcct gccacaccct tgttttgaga tctctctaa 420
ggagcggaga gtttaatagg caagaaggaa gggagaagac agaaggaga cgctccccg 480
tacggagaca gagggaaggg gggctccaaa gccgaaagag gaggctcccta cctgccacg 540
15 ataccagtca gcccttgcca gcatccagcc atgggggata tgaaaacccc agattttgat 600
gaccttctgg ctgcctttga catccagac ccaccagcc ttgatgcaa ggaggccatc 660
cagacaccca gtgaggagaa tgagagtccc ctcaaaccct caggcatatg tatggatgaa 720
agtgtgtcct tgtctcactc aggtacagcc ccgatgtgc cggccgtgag tgtcattgtc 780
aagaacacca gccgcagga gtcatttgaa gcggagaaag accacattac tcccagtctc 840
20 ctacacaatg gattccgggg ctccagatct cctccagatc ccacacaactg tgggaaattt 900
gatttacttt ttatgaatgg agacagtgc aggtgttcc ctggcaaaact ggagcctccc 960
aagtcagagc cattaccac ctccaaccag ttcagtccaa tctccagccc agaactgag 1020
gatcccatca aagataaacg atttgggata aagcccaaac actctgacag ttatttccca 1080
ccccctcttg ggtgcggggc tgtggggagg ccagtcctgg aggtcttggc taagtttccg 1140
25 gttccagagc tgcataatgt tgatcatttt tgtaagaaag aacccaagcc agaaccctcg 1200
cccttgggga gccagcagga acacgagcaa agtgggcaga acacagtgga acctcacaag 1260
gatccggatg ccactcgatt ctccggggaa gctttggagt tcaacagcca tcttagcaac 1320
agtattggag agtccaagg gcttgcgcgg gagcttggta cctgtctatc agtccccct 1380
aggcagcgtc taaagccagc tcattccaag ctgtcctctt gtgtggcagc ctgtgtggcc 1440
30 ttgcaggcca aaagagtggc tagtgtcact aaggaggatc agcctggcca cacaaggat 1500
ctctcagggc cactaaaga gatttctaaa ggtagcccca aaatgcccac gtcaccaaag 1560
agtccccgga gccctctgga ggccactaga aaaagtatca agccatcgga cagccctcgt 1620
agcatctgca gtgacagcag cagcaaggc tcaccgtctg tggctgccag ctccccacca 1680
gcaattccca aagtgagaat caaaaccatt aagacatcat caggggaaat caaacggact 1740
35 gtcacaagga tcttgccaga tctgatgat ccaagtaagt ccctgttgg gtcacctcta 1800
gggagcgcga ttgcagaggg cccagcgag atgcccaggg atgaggtgcc tgtggaagag 1860
cactttcctg aggcagggca aaattcaggg agccccagg gggccaggaa aggggacgag 1920
agcatgacaa aggcagtgta ctgctcact cccagctgca gttctgggcc cgggtccca 1980
aagggggctg cccagggctc acagacaggc aagaagcaac agagcacagc actgcaggca 2040
40 tccacctggt cccctgccaa cctcctgcc aaagccgtgc acttggccaa cctgaacctc 2100
gtccccaca gtgtgtgtgc atcagtgaac gccaaagtct cagtgcacaa acggagccag 2160
ccacagctta cacaatgtc ggtgccctg gtcccagg tgaaaaaggc tgcctcactg 2220
attgtagagg tcttcaacaa ggtccttcac agctccaacc ccgtgccct ctatgcgcca 2280
aatctcagcc cgcctgcgga cagcaggatc cacgtgccgg ccagtgggta ctgctgcctg 2340
45 gagtgtggag acgcatttgc cttagagaag agcctgagcc agcactatgg ccggcggagc 2400
gtccacattg aggtactgtg cacactgtgc tccaagacgc tgctcttctt caacaagtgc 2460
agcctgtccc ggcacgcccc tgaccacaag agcaaggggc tcgtcatgca gtgttccag 2520
ctgctggtga agcctatctc tgcggaccaaa atgttcgtgt cggccctgt gaactccacg 2580
gcaccagcag cccagcccc ttcatcctct cccaaacatg gcctcacttc gggcagtgcc 2640
50 agtccccctc ctccagcctt gccactctac ccagacctg tgaggctcat ccggtactca 2700
atcaagtgtc ttgaatgtca caagcagatg cgggactaca tggctcctgg tgcacatttc 2760
cagaggacaa cagaggagac agaggggctg acctgccagg tatgccagat gctgtgcc 2820
aaccagtga gtttctgtgc ccaccagcgg attcatgcac acaagteccc ctactgtgc 2880
ccggagtgtg gggctcctctg ccgctctgcc tacttccaga cccatgtaaa ggagaattgc 2940
55 ctgcactatg cccgcaaggt gggctacagg tgcattccact gtggtgtcgt ccacctgacc 3000
ttggccttgc tgaaaagcca catccaggag cgacactgcc aggttttcca caaatgtgca 3060
ttctgcccc tggccttcaa gactgccagc accacagtgc caccagcac 3120
cccacccagc cccacagacc ctcccagctc atttataagt gctcctgtga aatggtcttc 3180
aacaagaaga ggcacattca gcagcatttt taccagaatg tcagcaagac gcaggtgggc 3240
60 gtcttcaagt gccctgagtg cccactcttg ttcgtgcaga agccggagtt gatgcaaac 3300
gtcaagagca cccacggtgt tcccgaagt gtggacgagc tgtcaaacct ccagtcttca 3360
gcggacacat cctcaagccg cctggtctct cgagttccca ctgagccacc agccactagt 3420
gtggctgtc ggagcagctc cctgccttct ggccgctggg gtaggcctga agccaccgc 3480
aggttgaag ccaggccgcg gctgaggaa actggctgga cctgccagga gtgccaggag 3540

- 6 -

<400> 6
tttttttttt ttttttgtcc gacatcttga gacgaggtcg cggtgtctgc tgctattctc 60
cgagcttcgc aatgccgcct aaggacgaca agaagaagaa ggacgctgga aagtcggcca 120
agaaagacaa agacccagtg aacaaatccg ggggcaaggc caaaaagaag aagtggtcca 180
5 aaggcaaatg tcgggacaa gctcaataact tagtcttgtt tgacaaagct acctatgata 240
aactctgtaa ggaagtcccc aactataaac ttataacccc agctgtggtc tctgagagac 300
tgaagattcg aggtctccctg gccagggcag cccttcagga gctccttagt aaaggactta 360
tcaaactggt ttcaaagcac agagctcaag taatttacac cagaaatacc aagggtggag 420
atgctccagc tgctggtgaa gatgcatgaa taggtccaac cagctgtaca tttggaaaaa 480
10 taaaacttta ttaaatc 497

<210> 9
<211> 2056
15 <212> DNA
<213> Homo sapiens

<400> 9
20 ggaaccgagg ctgctggaca agaggggtgc ggtggatact gacctttgct ccggcctcgt 60
cgtgaagaca cagcgcatct ccccgctgta ggcttctccc acagaaccgg ttccgggcct 120
cagagcgtct ggtgagatgc tgttgccgct gctgctgctg ctaccatgt gctgggccgt 180
ggagggtcaag agggcccggg gcgtctccct caccatcat cacttctacg atgagtccaa 240
gcctttcacc tgcctggagc gttcggccac catccattt gatcaggtca acgatgacta 300
25 ttgcgactgc aaagatggct ctgacgagcc aggcacggct gcctgtccta atggcagctt 360
ccactgcacc aacactggct ataagccct gtatatcccc tccaaccggg tcaacgatgg 420
tgtttgtgac tgctgcgatg gaacagacga gtacaacagc ggcgtcatct gtgagaacac 480
ctgcaaagag aagggccgta aggagagaga gtccctgcag cagatggccg aggtcaccgg 540
cgaagggttc cgtctgaaga agatccctat tgaggactgg aagaaggcac gggaggagaa 600
gcagaaaaag ctcatgtgac tacaggctgg gaagaagtct ctggaagacc aggtggagat 660
30 gctgcggaca gtgaaggagg aagctgagaa gccagagaga gaggccaaag agcagcacca 720
gaagctgtgg gaagagcagc tggctgctgc caaggcccaa caggagcagg agctggcggc 780
tgatgccttc aaggagctgg atgatgacat ggacgggacg gtctcgggtg ctgagctgca 840
gactcaccgg gagctggaca cagatgggga tggggcgttg tcagaagcgg aagctcaggc 900
cctcctcagt ggggacacac agacagacgc cactctttc tacgaccgag tctgggccgc 960
35 catcagggac aagtaccggt ccgaggcact gcccaccgac cttccagcac cttctgcccc 1020
tgacttgacg gagcccaagg aggagcagcc gccagtcccc tcgtcgcccc cagaggagga 1080
ggaggaggag gaggaggagg aagaagaggc tgaagaagag gaggaggagg aggattccga 1140
ggaggcccca ccgccaactg caccgccgca gccggccagc cctgctgagg aagacaaaat 1200
gccgcccctac gacgagcaga cgcaggcctt catcgatgct gccaggagg cccgcaacaa 1260
40 gttcgaggag gccgagcggc cgctgaagga catggaggag tccatcagga acctggagca 1320
agagatttct tttgactttg gcccaacagg ggagtttctg tacctgtaca gccagtgtca 1380
cgagctcacc accaacgaat acgtctaccg cctctgcccc ttcaagcttg tctcgagaa 1440
acccaaactc gggggctctc ccaccagcct tggcacctgg ggctcatgga ttggccccga 1500
ccacgacaag ttcagtcca tgaagtatga gcaaggcacg ggctgctggc agggcccaaa 1560
45 ccgctccacc accgtgcgcc tctgtgcgg gaaagagacc atggtgacca gcaccacaga 1620
gcccagtcgc tgcgagtacc tcatggagct gatgacgcca gccgctgcc cggagccacc 1680
gcctgaagca cccaccgaag acgaccatga cgagctctag ctggatgggc gcagagaacc 1740
tcaagaaggc atgaagccag cccctgcagt gccgtccacc cgccccctg ggctgacctg 1800
tggtctgtgt gccctcctct gtggcgccag gacctttgtg gggcttcgtg cctctgctctg 1860
50 gggccccaggc ggggctggtc cacattccca ggccccaaac gcctccaaag atgggtaaag 1920
gagcttgccc tccctgggac cccacacctg gtgactcgcc ccaccacccc cagccctgtc 1980
cctgccaccc ctcctagtgg ggactagtga atgacttgac ctgtgacctc aatacaataa 2040
atgtgatccc ccaccc 2056

55 <210> 10
<211> 4577
<212> DNA
60 <213> Homo sapiens

<400> 10
gctacaatcc atctggtctc ctccagctcc ttctttctgc aacatgggga agaacaaact 60
ccttcaccca agtctggttc ttctcctctt ggtcctctg cccacagacg cctcagctctc 120
tggaaccg cagtataagg ttctgggtccc ctccctgctc cactactgaga cactgagaa 180

- 8 -

5 ctccttgaaa tacaatatc tcccagaaaa ggaagagttc ccctttgctt taggagtgc 4080
 gactctgcct caaacttgtg atgaacccaa agcccacacc agcttccaaa tctccctaag 4140
 tgtcagttac acagggagcc gctctgctc caacatggcg atcggtgatg tgaagatgg 4200
 cctctgcttc attccctcga agccaacagt gaaaatgctt gaaagatcta accatgtgag 4260
 ccggacagaa gtcagcagca accatgtctt gatttacctt gataaggtgt caaatcagac 4320
 actgagcttg ttcttcacgg ttctgcaaga tgtccagta agagatctca aaccagccat 4380
 agtgaaaagtc tatgattact acgagacgga tgagtttgca atcgctgagt acaatgctcc 4440
 ttgcagcaaa gatcttgaa atgcttgaag accacaaggc tgaaaagtgc tttgctggag 4500
 10 tctctgtctc tgagctccac agaagacacg tgtttttgta tctttaaaga cttgatgaat 4560
 aaacactttt tctggtc 4577

<210> 11

<211> 1651

15 <212> DNA

<213> Homo sapiens

<400> 11

20 ggagcccccg ccctgggatt cccaggtgtt ttcatttggg gatcagcact gaacacagaa 60
 gagtcatgac ggagtttggg ctgagctggg ttttccttgt tgctattttt aaaggtgtcc 120
 agtgtgaggt gcagctgggt gagtctgggg gagacttggg ccagcctggg gggctccctga 180
 gactctcctg tgcagcctct ggattcacct tcagtagtta tgctatgcac tgggtccgccc 240
 aggtccagg gaagggactg aaatatgttt caggtattag tagtaatggg cgtagaacat 300
 attatgcaaa ttctgtgaag ggcagattca ccatctccag agacaattcc aagaacacgt 360
 25 tgtatcttca aatgggcagc ctgagagctg aggacatggc tgtgtattac tgtgtgtccg 420
 ggggaatcta tgatagtagt ggtccctttg actactgggg ccagggaaacc ctggtcaccg 480
 tctcctcagc atccccgacc agccccaaagg tcttcccgct gagcctctgc agcaccacgc 540
 cagatgggaa cgtggctcctc gctgctctgg tccagggtct ctccccccag gagccactca 600
 gtgtgacctg gagcgaaagc ggacagggcg tgaccgccag aaacttccca cccagccagg 660
 30 atgctcccg ggacctgtac accacgagca gccagctgac cctgccggcc acacagtgc 720
 tagccggcaa gtccgtgaca tgccacgtga agcactacac gaatcccagc caggatgtga 780
 ctgtgccctg cccagtcccc tcaactccac ctaccccatc tccctcaact ccacctaccc 840
 catctccctc atgctgccac ccccgactgt cactgcaccg accggccctc gaggacctgc 900
 tcttaggttc agaagcgaac ctacgtgca cactgaccgg cctgagagat gcctcaggtg 960
 35 tcaccttcac ctggacgccc tcaagtggga agagcgtgt tcaaggacca cctgagcgtg 1020
 acctctgtgg ctgctacagc gtgtccagt tccgtccggg ctgtgccgag ccatggaacc 1080
 atgggaagac cttcacttgc actgtgcct accccgagtc caagaccggc ctaaccgcca 1140
 cctctcaaa atccggaaac acattccggc ccgaggtcca cctgtgccc ccgcccgcgg 1200
 aggagctggc cctgaacgag ctgggtgacgc tgacgtgcct ggacgcggc ttcagcccca 1260
 40 aggacgtgct gggtcgtgg ctgcagggtt cacaggagct gccccgagag aagtacctga 1320
 cttgggcac cggcaggag cccagccagg gcaccaccac cttcgtgtg accagcatac 1380
 tgcgcgtggc agccgaggac tggaaagaag gggacacctt ctctgcatg gtgggccacg 1440
 aggccctgcc gctggtcttc acacagaaga ccatcgaccg cttggcgggt aaaccacccc 1500
 atgtcaatgt gtctgtgttc atggcggagg tggacggcac ctgctactga gccgcccggc 1560
 45 tgtccccacc cctgaataaa ctccatgctc ccccaaaaaa aaaaaaaaaa aaaaaaaaaa 1620
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 1651

<210> 12

50 <211> 1120

<212> DNA

<213> Homo sapiens

<400> 12

55 acttcttcgc accaggggaag cccccaccac cagaacgcca agatgtccag caagcgggccc 60
 aaagccaagg ccaccaagaa gcggccacag cgggccacat ccaatgtctt cgcaatgttt 120
 gaccagtccc agatccagga gtttaaggag gctttcaaca tgattgacca gaaccgtgat 180
 ggcttcattg acaaggagga cctgcacgac atgctggcct cgtggggaa gaaccccaca 240
 gacgaatacc tggagggcac gatgagcgag gccccggggc catacaactt caccatgttc 300
 60 ctcaccatgt ttggggagaa gctgaacggc acggaccccg aggatgtgat tcgcaacgcc 360
 tttgctgct tcgacgagga atcctcaggt ttcatccatg aggaccacct ccggaagctg 420
 ctcaccacca tgggtgaccg cttcacagat gaggaagtgg acgagatgta ccgggaggca 480
 cccgttgata agaaaggcaa cttcaactac gtggagtcca cccgcaccc caaacatggc 540
 gccaaaggata aacacgacta ggccatcccc agccccctga caccacagccc ccgcccagtc 600

<210> 14

<211> 3612

<212> DNA

5 <213> Homo sapiens

<400> 14

10 aagcttaaaag tgaagacttg ctgaatttat ttcagcattt tactcctggc ctttttccat 60
aggaaggctc ctaatagtgt ctaactcctg aaagcagaaa gtgaccttca aaactgaccc 120
atcttccctg gaatctctct aagagctgag ggattagggg tcttagagat ttgtgtatct 180
ttgatggcac taaaatgaca tagagtttac atgtatacac ctggaaaagg tattaatagt 240
tatcagatgg ccctcaaaag tctagaacat ggtagctgtg cattgaggaa acattctgca 300
aaaaccattg tctcagaag aaacatccag ctctatatta attgtatgct cagacccatc 360
cctgctgggt aggtcagcat ttgtacttgg ttgcatatt ataccattgt gtgacatcaa 420
15 caatctgcaa gtacacattg ccgggctgcc tgtgcagaat ggcagctgct gggcctagag 480
tctagtgcag gtacttttga agggatcatt ctactgaat atgtctgttc tcttctatgt 540
actttctgac cctccagggt agctggatag agtggacctg aataagctaa acgagattga 600
aggcacccta aacaaagcca aagatgaaat gaaggtcagc gatcttgata ggaaagtgtc 660
tgacctggag aatgaagcca agaagcagga ggctgccatc atggactata accgagatat 720
20 cgaggagatc atgaaggaca ttgcgaatct ggaggacatc aggaagacct taccatctgg 780
ctgcttcaac acccctgcca ttgaaaagcc ctagtgtctt tagggctgga aggcagcatc 840
cctctgacag gggggcagtt gtgaggccac agagtgcctt gacacaaaga ttacattttt 900
cagaccccca ctctctgct gctgtccatc actgtccttt tgaaccagga aaagtcacag 960
agttaaaga gaagcaaat aaacatcctg aatcggaac aaagggtttt atctaataaa 1020
25 gtgtctcttc catcacgttg ctaccttacc cacacttccc tctgatttgc gtgaggacgt 1080
ggcatcctac ttacgtacgt ggcataacac atcgtgtgag cccatgtatg ctggggtaga 1140
gcaagttagc ctcccctgtc tcatcgatcc agcagaacct cctcagcttc agtactcttg 1200
ttctataag gaaaagtgtt gctactaaca gtacattgt gatggccagt atatccagtc 1260
catggataaa gaaaatgcat ctgcattctc tgccctctt ccttctaagc aaaaggaaat 1320
30 aaacatcctg tgccaaaggt attggtcatt tagaatgtcg gttagccatc atcagtgtct 1380
ttagctatta tgagtgtagg aacttgaccc atccgtgggt caggatgcaa ttatttataa 1440
aagtcccccag gtgaacatgg ctgaagattt ttctagtata ttaataattg actaggaaga 1500
tgaacttttt ttcatgcttt tgggcagctg ataattttaa tctggatggg cagcttgcat 1560
tcaccaatag accaaaagac atcttttgat attcttataa atggaactta cacagaagaa 1620
35 atagggatat gataaccact aaagttttgt ttccaaatc aaactaatc ttacagcttt 1680
tttattagtt agtcttgaa ctagtgttaa gtatctggca gagaacagtt aatccctaag 1740
gtcttgacaa aacagaagaa aaacaagcct cctcgtccta gtcttttcta gcaagggat 1800
aaaacttaga tggcagcttg tactgtcaga atcccggtga tccatttgtt cttctgttgg 1860
agagatgaga catttgaccc tttagctccag ttttctctg atgtttccat cttccagaat 1920
40 cctcaaaaa acattgtttg ccaaatcctg gtggcaata cttgcactca gtatttcaca 1980
cagctgcaaa cgctatcgag ttctgcact ttgtgattta aatccactct aaaccttccc 2040
tctaagtgtg gagggagac ccttacgtgg agtttcctag tgggcttctc aacttttgat 2100
cctcagctct gtggttttaa gaccacagtg tgacagtccc ctgccacaca ccccttctct 2160
cctaccaacc cacccttgag attcatatat agcctttaac actatgcaac ttgtactttt 2220
45 gcgtagcagg ggctggggg gggggaaaga aacctattat catggacaca ctgggtctat 2280
taattatttc aaatttatat tttgtgtgta atgttttgtg tttgttttat ccatgctata 2340
gaacaaggaa tttatgtaga tatacttagt cctatttcta gaatgacact ctgttcaact 2400
tgctcaattt ttctcttcca ctggcacaag tatctgaata cctccttccc tcccttctag 2460
agtctcttgg attgtactcc aaagaattgt gccttgtgtt tgcagcatct ccattctcta 2520
50 aattaatata attgtcttcc tccacacca gccacgtaaa gaggttaact gggctctctt 2580
ccattgcagt cctgatgac ctaacctgca gcacgggtgt tttacaatgt tccagagcag 2640
gaacgccagg ttgacaagct atggtaggat taggaaagtt tgctgaagag gatcttgac 2700
gccacagtgg gactagccag gaatgaggga gaaatgccct ttttggcaat tgttggagct 2760
ggataggtaa gttttataag ggagtacatt ttgactgagc acttagggca tcaggaacag 2820
55 tgctacttac tgggtggtag actgggagag gtggtgtaac ttagtctctg atgatccac 2880
ttctgtttc catctgctg ggatatacca gagtttacca caagtgtttt gacgatatac 2940
tcttgagctt tcaactctgt ggcttctccc aggcctcttc tactatggca ggagatgtgg 3000
tgtgtgttgg caaagttttc acgtcatcgt ttctggcta gttcatttca ttaagtggct 3060
acatcctaac atatgcattg gtcaagggtg cagcaaggag actgaagatt gactgccaag 3120
60 ctagtgttgg tgaagttcac tccagcaagt ctccaggccac aatgggggtg tttgggttgg 3180
tttctttta actttctttt tgttatttgc ttttctctc cacctgtgtg gtatattttt 3240
taagcagaat tttatttttt aaaataaaa gttctttaca agatgatacc ttaattacac 3300
tcccgaaca cagccattat tttattgtct agctccagtt atctgtattt tatgtaattg 3360
aatgacagg atggctgctg cagaatgctg gttgacacag ggattattat actgctattt 3420

ccggaggtga agtcgggtgcc acttctgccc tggcccccaa gatcggcccc ctgggtctgt 180
 ctccaaaaaa agttgggtgat gacattgcca aggcaacggg tgactggaag ggcttgagga 240
 ttacagtga actgaccatt cagaacagac aggccagat tgaggtggtg ccttctgcct 300
 ctgccctgat catcaaagcc ctcaaggaa caccaagaga cagaaagaaa cagaaaaaca 360
 5 ttaaacaacag tgggaatatc acttttgatg agattgtcaa cattgtctga cagatgcggc 420
 accgatcctt agccagagaa ctctctggaa ccattaaaga gatcctgggg actgcccagt 480
 cagtgggctg taatgttgat ggccgccatc ctcatgacat catcgatgac atcaacagtg 540
 gtgctgtgga atgcccagcc agttaagcac aaaggaaaac atttcaataa aggatcattt 600
 gacaactggg ga 612
 10
 <210> 17
 <211> 1100
 <212> DNA
 15 <213> Homo sapiens
 <400> 17
 ctgcagtatt tagcatgccc caccatctg caaggcattc tggatagtgt caaaacagcc 60
 ggaaatcaag tccgtttatc tcaaaacttta gcattttggg aataaatgat atttgctatg 120
 20 ctggttaaat tagattttag ttaaaatttc tgctgaagct ctagtacgat aagcaacttg 180
 acctaatgtt aaagttaga tttccttcag gtttatatag ctgtgcgcc gcctgggtac 240
 ctccgagtgt agggcgatct ggctgcgaca tctgtcacc cttgtatcgc cagggttgat 300
 ctggctgatc tggctggcta ggctgggtgtc cccttctctc ctcccgctc catgtgcgtc 360
 cctccggaag ctgcgcgctc ggtcgaagag gacgaccatc cccgatagag gaggaccggg 420
 25 ctccggtcaa gggatatacg gtatgtgcgc tccctgtcta gaacctcaa acaagctctc 480
 aaggtccatt tgtaggagaa cgtagggtag tcaagcttcc aagactccag acacatccaa 540
 atgaggcgct gcatgtggca gtctgccttt ctttgaccc attaccatc taaggaccgg 600
 tcttcgggtca aggggtatacg agtagctgcg ctccctgtct agaacctcca acaagctctc 660
 caaggtccat ttgtaggaga acgtagggta gtcaagcttc caagactcca gacacatcca 720
 30 aatgaggcgc tgcattgtggc agtctgcctt tcttttgacc cattaccat ctaagttaga 780
 tgctttttta aatgtttttt aatttttaaa tttttaattt ttttcattat ttatttttta 840
 tttttgagac ggaatctcggc tcactgtaac ctccacctcc cgggttcaag cgattctcct 900
 gactcagcct cccgagtagc tgggattaca ggcgcgcgcc accatgccc gctaattttt 960
 gtatttttag taaagacggg ttttcgcat gttagaccag gtggtcttga actcctgacg 1020
 35 tcaggtgatc ctctcgcctc gtgctgggat tgcaggcgct agccaccgct cccggccctt 1080
 agatgctttt taatcagcaa 1100
 <210> 18
 40 <211> 4747
 <212> DNA
 <213> Homo sapiens
 <400> 18
 45 taggaatgaa ccatccctgg gagtatgtgg aaacaacgga ggagctctga cttcccaact 60
 gtcccatctt atgggcgaag gaactgctcc tgacttcagt gggttaagggc agaattgaaa 120
 ataattctgg aggaagataa gaatgattcc tgcgcgactg caccgggact acaaagggtc 180
 tgtcctgctg ggaatcctcc tggggactct gtgggagacc ggatgcaccc agatacgcta 240
 ttcagttccg gaagagctgg agaaaggctc taggggtggg gacatctcca gggacctggg 300
 50 gctggagccc cgggagctcg cggagcgcg agtccgcac atcccagag gtaggacgca 360
 gcttttcgcc ctgaatccgc gcagcgccag cttggtcacg gcgggcagga tagaccggga 420
 ggagctctgt atggggggcca tcaagtgtca attaaatcta gacattctga tggaggataa 480
 agtgaaaata tatggagtag aagtagaagt aaggacatt aacgacaatg cgccttactt 540
 tctgtaaagt gaattagaaa taaaaattag tgaatatgca gccactgaga tgcggttccc 600
 55 tctacccac gcctgggac cggatatcgg gaagaactct ctgcagagct acgagctcag 660
 cccgaacact cacttctccc tcatcgtgca aaatggagcc gacggtagta agtaccgccg 720
 attggtgctg aaacgcgccc tggaccgca agaaaaggct gtcaccacc tggctcttac 780
 ggctccgac gggggcgacc cgggtgcgac aggcaccgag cgcatccgag tgatggttct 840
 ggatgcgaac gacaacgcac cagcgtttgc tcagcccgag taccgcgca gcgttcggga 900
 60 gaatctggcc ttgggcacgc agctgcttgt agtcaacgct accgaccctg acgaaggagt 960
 caatgcggaa gtgaggtatt ccttcgggta tgtggacgac aaggcgggcc aagttttcaa 1020
 actagattgt aattcagga caatatcaac aataggggag ttggaccagc aggagtcagg 1080
 attctaccag atggaagtgc aagcaatgga taatgcagga tattctgcgc gagccaaagt 1140
 cctgatcact gttctggacg tgaacgacaa tgccccagaa gtggtcctca cctctctcgc 1200

<212> DNA

<213> Homo sapiens

<400> 19

5 gaattccgca gcccgagtg tggtagcag ctcggcaagc gctgcccagg tcctggggtg 60
gtggcagcca gcgggagcag gaaaggaagc atgttcccag gctgcccagg cctctgggtc 120
ctggtgggtc tgggacccag ctgggtagcc tgggggagcc aagggacaga agcggcacag 180
ctaaggcagt tctacgtggc tgctcagggc atcagttgga gctaccgacc tgagcccaca 240
aactcaagtt tgaatctttc tgtaactttc ttttaagaaa ttgtctacag agagtatgaa 300
10 ccatatttta agaaagaaaa accacaatct accatttcag gacttcttgg gcctacttta 360
tatgtgaaag tcggagacat cataaaagtt cactttaaaa ataaggcaga taagcccttg 420
agcatccatc ctcaaggaaat taggtacagt aaattatcag aaggtgcttc ttaccttgac 480
cacacattcc ctgaggagaa gatggacgac gctgtggctc caggccgaga atacacctat 540
gaatggagta tcagtgaaga cagtggaccc accatgatg accctccatg cctcacacac 600
15 atctattact cccatgaaaa tctgatcgag gatttcaact cggggctgat tgggcccctg 660
cttatctgta aaaaaggagc cctaactgag ggtgggacac agaagacgtt tgacaagcaa 720
atcgtgctac tatttgctgt gtttgatgaa agcaagagct ggagccagtc atcatcccta 780
atgtacacag tcaatggata tgtgaatggg acaatgccag atataacagt ttgtgcccat 840
gaccacatca gctggcatct gctgggaatg agctcggggc cagaattatt cctcattcat 900
20 ttcaacggcc aggtcctgga gcagaacat cataaggtct cagccatcac cctgtcagt 960
gtacatccca ctaccgcaaa tatgactgtg ggcccagagg gaaagtggat catatcttct 1020
ctcaccctcaa aacattttga agctgggatg caggcttaca ttgacattaa aaactgccca 1080
aagaaaacca ggaatcttaa gaaaataact cgtgagcaga ggcggcacat gaagaggtgg 1140
gaatacttca ttgctgcaga ggaagtcat tgggactatg cactgtaat accagcgaat 1200
25 atggacaaaa aatacaggtc tcagcatttg gataatttct caaaccacat tggaaaacat 1260
tataagaaag ttatgtacac acagtacgaa gatgagtcct tcaccaaaca tacagtgaat 1320
cccaatatga aagaagatgg gattttgggt cctattatca gagcccaggt cagagacaca 1380
ctcaaaatcg ttttcaaaaa tatggccagc cgcccctata gcattttacc tcatggagt 1440
accttctcgc cttatgaaga tgaagtcaac tcttctttca cctcaggcag gaacaacacc 1500
30 atgatcagag cagttcaacc aggggaaacc tatacttata agtggacat cttagagttt 1560
gatgaaccca cagaaaatga tggccagtgc ttaacaagac cactactacag tgacgtggac 1620
atcatgagag acatcgccctc tgggctaata ggactacttc taatctgtaa gacgagatcc 1680
ctggacaggg gaggaatata gagggcagca gacatcgaaac agcaggctgt gtttgcgtg 1740
ttgatgaga acaaaaagctg gtaccttgag gacaacatca acaagttttg tgaaaatcct 1800
35 gatgaggtga aacgtgatga ccccaagttt tatgaatcaa acatcatgag cactatcaat 1860
ggctatgtgc ctgagagcat aactactctt ggattctgct ttgatgacac tgtccagtgg 1920
cacttctgta tgttggggac ccagaatgaa attttgacca tccacttcac tgggactcca 1980
ttcatctatg gaaagaggca tgaggacacc ttgaccctct tcccctatgc tggagaatct 2040
gtgacggtca caatggataa tgttggaaact tggatgttaa cttccatgaa ttctagtcca 2100
40 agaagcaaaa agctgaggct gaaattcagg gatgttaaat gtatcccaga tgatgatgaa 2160
gactcatatg agatttttga acctccagaa tctacagtea tggctacacg gaaaatgcat 2220
gatcgtttag aacctgaaga tgaagagagt gatgctgact atgattacca gaacagactg 2280
gtcgagcat taggaattag gtcattccga aactcatcat tgaaccagga agaagaagag 2340
ttcaatctta ctgccctagc tctggagaat ggcactgaat tcgtttcttc gaacacagat 2400
45 ataattgttg gttcaaatla ttcttcccca agtaatatla gtaagttcac tgtcaataac 2460
cttgacagaa ctacagaaagc cccttctcac caacaagcca ccacagctgg tccccactg 2520
agacacctca ttggcaagaa ctacgttctc aattcttcca cagcagagca ttccagccca 2580
tattctgaag acctataga ggtacctcta cagccagatg tcacagggat acgtctactt 2640
tcacttggtg ctggagaatt cagaagtcaa gaacatgcta agcgtaaagg acctaaaggt 2700
50 gaaagagatc aagcagcaaa gcacaggttc tcctggatga aattactagc acataaagtt 2760
gggagacacc taagccaaga cactggttct ccttcgggaa tgaggccctg ggaggacctt 2820
cctagccaag aactggttc tcttccaga atgaggccct gggaggaccc tctagtgat 2880
ctgttactct taaaacaaag taactcatct aagattttgg ttgggagatg gcatttggct 2940
tctgagaaag gtacttatga aataatccaa gatactgatg aagacacagc tgttaacaat 3000
55 tggctgatca gccccagaa tgctcacgt gcttggggag aaagcaccct tcttgccaac 3060
aagcctggaa agcagagtgg ccacccaag tttcttagag ttagacataa atctctacaa 3120
gtaagacagg atggaggaaa gactagactg aagaaaagcc agtttctcat taagacacga 3180
aaaaagaaaa aagagaaagc cacacacat gctcctttat ctccaggagc ctttaccct 3240
ctaagaagtg aagcctacaa cacattttca gaaagaagac ttaagcatc gttgtgctt 3300
60 cataaatcca atgaacatc tcttccaca gacctcaatc agacattgcc ctctatggat 3360
tttggctgga tagctcact tctgacct aatcagaatt cctcaaatga cactggtcag 3420
gcaagctgtc ctccaggtct ttatcagaca gtgccccag aggaacacta tcaaacatc 3480
ccattcaag acctgatca aatgcactct acttcagacc ccagtcacag atcctcttct 3540
ccagagctca gtgaaatgct tgagtatgac cgaagtcaca agtccttccc cacagatata 3600

ccgcccgtt ctactccgcc gcgggggtcg cagcggtgc cgcgcgtcc tcgagttcc 60
agcgtgagga ggaggctgag ggaggagagg cgcacgtgt tcgaggcggg gaccgagggg 120
gagccccgcg cgcggcgctg ctcatgtcta tggacagtgc tatccacctg tggcagttcc 180
ttcttcagct cctgcagaag cctcagaaca agcacatgat ctgttggacc tctaagtatg 240
5 ggcagtttaa gcttttgcag gcagaagagg tggctcgtct ctgggggatt cgcaagaaca 300
agcctaacat gaattatgac aaactcagcc gagccctcag atactattat gtaagaata 360
tcatcaaaaa agtgaatggt cagaagtgtg tgtacaagt tgtctcttat ccagagattt 420
tgaacatgga tccaatgaca gtgggcagga ttgagggtga ctgtgaaagt ttaaacttca 480
gtgaagtcag cagcagttcc aaagatgtgg agaattggagg gaaagataaa ccacctcagc 540
10 ctgggtccaa gacctctagc cgcaatgact acatacactc tggcttatat tcttcattta 600
ctctcaactc tttgaactcc tccaatgtaa agcttttcaa attgataaag actgagaatc 660
cagccgagaa actggcagag aaaaaatctc ctccaggagcc cacaccatct gtcatacaat 720
ttgtcacgac accttccaaa aagccaccag ttgaacctgt tgctgccacc atttcaattg 780
gcccagat tctctcatct tcagaagaaa ctatccaagc tttggagaca ttggtttccc 840
15 caaaactgcc ttccttggaa gcccacact ctgcctctaa cgtaatgact gcttttgcca 900
ccacaccacc catttctgccc ataccctctc tgcaggaaacc tcccagaaca ccttcaccac 960
cactgagttc tcaccagagc atcgacacag acattgattc agtggcttct cagccaatgg 1020
aaacttcaga gaatttgtct ctggagccta aagaccagga ttcagtcttg ctagaaaagg 1080
acaaagttaa taattcatca agatccaaag aacccaaagg gttaggactg gcaccacccc 1140
20 ttgtgatcac gagcagtgat ccaagcccac tgggaatact gagcccatct ctccttacag 1200
ctctctttac accagcattt tttcacaga caccatcat actgactcca agcccttgc 1260
tctccagtat ccaacttctg agtactctca gtctgttgc tcccctaagt ccagccagac 1320
tgcaaggtgc taacacactc ttcagtttc ctctgtact gaacagtcac gggccattca 1380
ctctgtctgg gctggatgga ccttccaccc ctggcccatt tccccagac ctacagaaga 1440
25 cataacctat gcacttggg aatgagagaa ccgaggaacg aagaaacaga cattcaacat 1500
gattgcattt gaagtgaaga atgtatgatt ctacaatgct gataatagac tattgtgatt 1560
tttgccattc cccattgaaa acatctttt aggattctct ttgaatagga ctcaagtgg 1620
actatagta taaaaatgcc ttaattggag tctaaactcc acctccctct gtcttttct 1680
tttcttttc tttccttctc tcttttctc tctccttta aaaatatctt gagctttgtg 1740
30 ctgaagaagt ttttgggtgg ctttagtgac tgtgtttgc aaaagcaatt aagaacaaag 1800
ttactccttc tggctattgg gacccttgg ccaggaaaaa ttatgcttag aatctattat 1860
ttaaagaagt atttgtgaaa tgaaaaaaa aaaaaaaa aaaaaaaa 1920
aaaaaaaaaaa aaa 1933

35 <210> 21
<211> 2215
<212> DNA
<213> Homo sapiens

40 <400> 21
actcagtaga ccgccactgg ctgtgcacgt tatggggttt ccacctaggg ctccgacctga 60
ggcttgaac actccgtttt cccccagtc acaggggcag tcttgcctct cgcagctggg 120
tcgcggtgtc tctcaaaggc cccctctac aggggcttcg tgaggcccg gcccacaggg 180
45 cgtcgggtcc cggaagtga gtctccaga ggggcccga gtggcagtg agggaggga 240
gatggcggag gtgggggaga taatcgaggg ctgccccta cccgtgtgc ggcggaacca 300
ggacaacgaa gatgagtgcc cctggccga gatcctgagc gtgaaggaca tcagtggccg 360
gaagcttttc tacgtccatt acattgactt caacaaacgt ctggatgaat gggtagcga 420
tgagcggctg gacctaaaga agatccagtt ccccaagaaa gaggccaaga cccccctaa 480
50 gaacggactt cctgggtccc gtcttgctc tccagagaga gagggtcccg cctcggcgca 540
ggccagcggg aagaccttgc caatcccggc ccagatcaca ctccgttca acctgcccaa 600
ggagcgggag gccattcccg gtggcgagcc tgaccagccg ctctcctcca gctcctgct 660
gcagcccaac caccgctcaa cgaaacggaa ggtggagggt gtttaccag caactccagt 720
gcccagcgag acagccccgg cctcgggttt tccccagaat ggagccgccc gtaggcgagt 780
55 ggcagcccag ccaggacgga agcgaaaatc gaattgtttg ggcactgatg aggactccca 840
ggacagctct gatggaatac cgtcagcacc acgcatgact ggcagcctgg tgtctgatcg 900
aagccacgac gacatcgtca cccggatgaa gaacattgag tgcattgagc tgggcccgca 960
ccgctcaag ccgtgtgact tctccccgta cccacaggaa ctccaccat tgctgtcct 1020
ctacctgtgc gaggttctgc tcaagtacgg ccgtagtctc aagtgtcttc agcgtcattt 1080
60 gaccaagtgt gacctacgac atcctccagg caatgagatt taccgcaagg gcaccatctc 1140
cttctttgag attgatggac gtaagaacaa gagttatctc cagaacctgt gtcttttggc 1200
caagtgttc cttgaccata agacactgta ctatgacaca gacccttcc tctctacgt 1260
catgacagag tatgactgta agggcttcca catcgtgggc tacttctcca aggagaaaga 1320
atcaacggaa gactacaatg tggctgcat cctaaccctg cctccctacc agcgcgggg 1380

- 18 -

ggtctctggtg gaacacagcc agacatcctg atggggaggc agagccagga agctaagcca 1320
 gggcccagct gcgtccaacc cagcccccca cctcagggtc ctgaccccag ctcatgccc 1380
 catcaccgcc tctccctggc tccccagggt gcagggtggc gcaaggccc gcccccatca 1440
 catgttccct tggcctcaga gctgcccctg ctctcccacc acagccaccc agaggcacc 1500
 5 catgaagctt ttttctcgtt cactcccaaa ccaagtgtc caaagctcca gtcctaggag 1560
 aacagtccct gggtcagcag ccaggaggcg gtccataaga atggggacag tgggctctgc 1620
 cagggtgccc gcacctgtcc agaacaacat gttctgttcc tctctctcat gcatttccag 1680
 ccttg 1685

10 <210> 24
 <211> 6428
 <212> DNA
 <213> Homo sapiens

15 <400> 24
 gctagtggaa gttactgccg cgcaccgag tccggaccgg agactttggg gcctaactag 60
 tgaatggtag tgtctagaaa gggatgtgcc cttcaagaga gaggtgcaa tgtccaaccg 120
 gcctaataac aatccagggg ggtcactgcy acgttcacag aggaacactg ccggggccca 180
 20 accacaagac gactcaatag gaggaagaag ctgcagttca tcatctgctg tgatagtccc 240
 acaaccagag gatccagaca gagccaatac ttcagaaaga caaaaaacgg ggcaggtgcc 300
 taagaaagac aattctcagag gagtgaagcg cagtgcctagt ccagactaca acaggacca 360
 ttctcctagc tctgcaaaaa aacaaaagc acttcagcat actgaatctc cctcagaaac 420
 aaataagcca catagtaagt caaagaagag acatttagac caggagcaac aactgaaatc 480
 25 tgcacaatca ccatcaacaa gcaaggctca taccagggaag agtggggcca ctggcggttc 540
 acggagtcag aaaagaaaaa ggacagagag ttcttgtgta aagagtggct ccgggtctga 600
 atcaactggt gcagaagaga gatctgcgaa acctaccaag ctggcttcaa aatcagccac 660
 ctcagccaaa gctgggtgta gcaccatcac tgattcttct tctgctgctc ctacttctc 720
 ctgctcttct gctgtagcct cggcctctc cactgtacca ccaggtgcca gagtgaaca 780
 30 aggaaaagat cagaacaagg ccaggcgttc ccgttcagcg tccagtccca gccccagaag 840
 aagtagcagg gaaaaggaac agagtaaaac tgggtggctct tcaaaatttg attgggctgc 900
 tcgtttcagc cctaaagtta gccttctctaa acaaaaactg tctcttccag ggtcttctaa 960
 gtcagagaca tcaaaaacctg gccttctgag attacaggcc aaattagcaa gtttaagaaa 1020
 atctacgaag aaacgcagtg agtctccacc tgctgagctc ccagtttga ggcggagcac 1080
 35 acgcaaaaag accacgggct cctgtgctag taccagtcgg cgaggctctg gcctgggcaa 1140
 aagaggagca gctgaagctc gtcgacagga gaaaatggca gacctgaaa gcaaccagga 1200
 ggcagtaaat tcttcaagctg ctccgacaga tgaagctccc caaggagctg caggggctgt 1260
 tggcatgacc acctctgggg agagtgaatc agatgattcc gagatgggac gtttgcaagc 1320
 tttgttagag gcaaggggct tccccctca cctatttggc cctcttggc ctcggtatgc 1380
 40 acagcttttc catagaacaa ttggaagtgg agctagtctt aaggcccagc agctaactaca 1440
 aggattgcaa gccagtgatg aaagtcaaca gcttcaggca gttattgaga tgtgtcagtt 1500
 actggtcatg ggaaatgagg agacactggg aggggttccct gtcaagagtg ttgttccagc 1560
 tttgattacg ttacttcaga tggagcacia ttttgatatt atgaaccatg cttgtcagac 1620
 cttaacatac atgatggaag cacttctctg atcttctgct gttgtagtag atgctattcc 1680
 45 tgtcttttta gaaaagctgc aagtatttca gtgtattgat gtggcagagc aggccttgac 1740
 tgccctggag atgttgctac ggagacatag taaagccatt ctacaggcgg gtggtttggc 1800
 agactgcttg ctgtacctag aattcttccag cataaatgcc caaagaaatg cattagcaat 1860
 tgcagctaatt tgcgtccaga gtatcacgcc agatgaattt cattttgtgg cagattcact 1920
 cccattgcta acccaaaagg taacacatca ggataaaaag tcagtagaaa gcacttgcct 1980
 50 ttgttttgca cgcctagtgg acaacttcca gcatgaggag aatttactcc agcaggttgc 2040
 ttccaaagat ctgcttcaaa atgttcaaca gctgttggta gtgactccac ccattttaag 2100
 ttctgggatg tttataatgg tggttcgcat gtttctctg atgtgttcca actgtccaac 2160
 tttagctggt caacttatga aacaaaacat tgcagaaacg cttcactttc tctgtgtgg 2220
 tgccctcaat ggaagtgtgc aggaacagat tgatcttgtt ccacgaagcc ctcaagagtt 2280
 55 gtatgaactg acatctctga tttgtgaact tatgccatgt ttacaaaag agggcatttt 2340
 tgcagttgat accatgttga agaagggaaa tgcacagaac acagatgggt cgatatggca 2400
 gtggcgtgat gatcggggcc tctggcatcc atataacagg attgacagcc ggatcattga 2460
 gcaaatcaat gaggcacagc gaacagcacg tgccattcag agaaaacctc acccgtagc 2520
 caatagtaac actagtggat attcagagtc aaagaaggat gatgctcgag cacagcttat 2580
 60 gaaagaggat ccggaactgg ctaagtcttt tattaagaca ttatttgggt ttctttatga 2640
 agtgatatgt tcttcagcag gacctgcggg cagacataag tgcccttagag caattcttag 2700
 gataatttat tttgcggatg ctgaacttct gaaggatgtt ctgaaaaatc atgctgttcc 2760
 aagtcacatt gcttccatgc tgtcaagcca agacctgaag atagtagtgg gagcacttca 2820
 gatggcagaa attttaatgc agaagttacc tgatattttt agtgtttact tcagaagaga 2880

- 20 -

<212> DNA

<213> Homo sapiens

<400> 25

5 gcacgaggcc agccttacgg gcccgaaacc tcgtgtgaag ggtgcagtac ctaagccgga 60
gcggggtaga ggcggggccg caccgccctt tgacctccag tgcgcggcgc ctcaagatca 120
gacatggccc agaacttgaa ggacttgccg ggacggctgc ccgcccggcc ccggggcatg 180
ggcacggccc tgaagctgtt gctgggggccc ggccgctggt cctacggtgt gcgcgaatct 240
gtgttcaccg tggaaaggcg gcacagagcc atcttcttca atcggatcgg tggagtgcag 300
10 caggacacta tcctggccga gggccttcac ttcaggatcc cttgggtcca gtacccatt 360
atctatgaca ttcggggccg acctcgaaaa atctcctccc ctacaggctc caaagacct 420
cagatggtga atatctccct gcgagtgttg tctcgacca atgctcagga gcttcctagc 480
atgtaccagc gcctagggct ggactacgag gaacgagtgt tgcggtccat tgttaatgag 540
gtgtccaaga gtgtgggtgc caagttcaat gcctcacagc tgatcaccca gcggggccag 600
15 gtatccctgt tgatccgccc ggagctgaca gagagggccca aggacttcag cctcatcctg 660
gatgatgtgg ccacacaga gctgagcttt agccgagagt acacagctgc tgtagaagcc 720
aaacaagtgg ccacgacgga ggcggcgccg gcccaattct tggtagaaaa agcaagcag 780
gaacagcggc agaaaattgt gcaggccgag ggtgaggccg aggtgcca gatgcttga 840
gaagcactga gcaagaacct tggctacatc aaacttcgca agattcgagc agcccagaat 900
20 atctccaaga cgatcgccac atcacagaat cgtatctatc tcacagctga caaccttgtg 960
ctgaacctac aggatgaaag ttaccaccag ggaagtgaca gcctcatcaa gggtaagaaa 1020
tgagcctagt caccaagaac tccaccccca gaggaagtgg atctgcttct ccagtttttg 1080
aggagccagc cagggggtcca gcacagccct acccgcccc agtatcatgc gatgggtccc 1140
cacaccgtt ccctgaacct ctcttggtt aagggaagact gaagactagc ccttttctg 1200
25 ggggaattact ttcctctcc ctgtgttaac tggggctgtt ggggacagtg cgtgatttct 1260
cagtgatctc ctacagtgtt gttccctccc tcaaggctgg gaggagataa acaccaacct 1320
aggaattctc aataaatttt tattacttaa cctgaaaaaa aaaaaaaaaa 1370

30 <210> 26

<211> 642

<212> DNA

<213> Homo sapiens

35 <400> 26

ccgcgcgtca cgtgacccca gcgcctactt gggctgagga gccgcgcgt cccctcgccg 60
agtcctctcg ccagattccc tccgtcgccg ccaagatgat gtgcggggcg ccctccgcca 120
cgcagccggc caccgcccag acccagcaca tcgcccacca ggtgaggtcc cagcttgaag 180
agaaagaaaa caagaagtcc cctgtgttta aggcggtgtc attcaagagc caggtggtcg 240
40 cggggacaaa ctacttcac aaggtgcacg tcggcgacga ggacttcgta cacctgcgag 300
tgttccaatc tctccctcat gaaaacaagc ccttgacctt atctaactac cagaccaaca 360
aagccaagca tgatgagctg acctatttct gatcctgact ttggacaagg cccttcagcc 420
agaagactga caaagtcac ctccgtctac cagagcgtgc acttgtgatc ctaaaataag 480
cttcatctcc gctgtgccc tggggtgga ggggcaggat tctgcagctg cttttgcatt 540
45 tctcttccca aatttcattg tgttgatttc tttccttccc aatagggtgat cttaattact 600
ttcagaatat tttcaaaata gatattttt taaatcctt ac 642